

A comparative study of cardiovascular manifestations in treatment naïve versus “on-treatment” patients with retroviral disease

Anugraha Durairaj, Hrishikeshshiva Kumar, Madhu G*

Post Graduate , Department of Medicine , BMCRI , Bangalore -560002, Karnataka , India

Received: 11-03-2021 / Revised: 12-04-2021 / Accepted: 07-05-2021

Abstract

Background: ART increases the longevity of the patients infected with HIV so mortality due to Opportunistic infections is reducing and late complications like cardiovascular manifestations are on its rise. Cardiac involvement can be over looked in HIV positive patients because symptoms of breathlessness, fatigue and poor exercise tolerance are ascribed to other conditions associated with HIV patients. **Objectives:** Primary: To evaluate cardiac manifestations and determine type of cardiac involvement in both ART naïve patients and patients on ART and correlate with CD4+ counts. **Methods:** 200 HIV infected patients presenting in OPD and inpatients are included. Information is collected and detailed history is taken using pre-formed proforma. At the time of admission or follow up, steps are taken to send for all the investigations and detailed clinical examination of the patient done, focusing more on cardiovascular system. **Results:** 41% of the patients with normal cardiac function were NOT ON ART and 62.7% of the patients with cardiac dysfunction were HIV naïve and it was found to be statistically significant (p value 0.003). Tricuspid regurgitation (42.1% vs 57.9%, p value 0.035) and IHD (0% vs 100% p value 0.049) found to have significant association in patients who were ON ART. Systolic dysfunction (p value 0.048) and IHD (p value 0.019) were both significantly associated with the low CD4+ counts in patients NOT ON ART. **Conclusion:** There was significant association of ART on the occurrence of cardiac dysfunction. Cardiac dysfunction is noted with low CD4+ Counts, it is therefore important to undergo 2DECHO at diagnosis and at regular intervals in all HIV infected patients and to initiate early treatment for the same.

Keywords: ART, HIV

This is an Open Access article that uses a fund-ing model which does not charge readers or their institutions for access and distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>) and the Budapest Open Access Initiative (<http://www.budapestopenaccessinitiative.org/read>), which permit unrestricted use, distribution, and reproduction in any medium, provided original work is properly credited.

Introduction

HIV infection is a major health problem in the entire world including India and globally 36.7 million [34.0 million–39.8 million] people were living with HIV at the end of 2015[1]. As per the NACO, India HIV Estimation 2015 report, National adult (15-49) years HIV prevalence in India is estimated at 0.26% in 2015 with 0.30% among males and 0.22% among females. The total number of people living with HIV in India is estimated at 21.17 lakhs (17.11 lakhs-26.49 lakhs) in 2015[2]. The prevalence of cardiac involvement in AIDS patients have been reported to range between 28% and 73%[3]. Cardiac involvement in AIDS/HIV infected persons may be attributed to virus itself, the effects of anti-retroviral medications; or altered immune mechanisms associated with the infection[4]. From the beginning of AIDS epidemic, cardiac involvement was recognized at autopsy and later by non-invasive techniques such as electrocardiogram. Patients with HIV infection can have variety of cardiovascular manifestations. The most common are pericarditis, pericardial effusion, pulmonary vascular diseases, myocarditis, cardiomyopathy and increased incidence of coronary artery disease, malignant neoplasm, coronary artery disease and drug related cardio toxicity[5]. The prevalence of heart muscle disease is around 15% and has 3 forms global left ventricular dysfunction, isolated right ventricular dilatation and borderline left ventricular dysfunction. Dilated cardiomyopathy is an independent adverse prognostic factor. It is strongly associated with very low CD4 cell count. Present study is undertaken to determine type of cardiac involvement in both ART

naïve patients and patients on ART and correlate pattern of cardiac involvement with CD4+ counts.

Objectives

Primary objective: To evaluate cardiac manifestations and determine type of cardiac involvement in both ART naïve patients and patients on ART and correlate pattern of cardiac involvement with CD4+ counts.

Secondary objective: To evaluate cardiac manifestations in patients having opportunistic infections.

Materials and methods

Source of data: The study would be conducted in patients with HIV treatment naïve and on ART admitted in Victoria, Bowring and Lady Curzon hospitals which are attached to Bangalore Medical College & Research Institution during the study period of January 2018-April 2020.

Methods of collection of data (including sampling procedure, if any)

A. Study design : Cross sectional study

B. Study period : January 2018-April 2020

C. Place of study: Victoria Hospital and Bowring and Lady Curzon Hospital, Bangalore Medical College & Research Institution , Bangalore

D. Sample size: 200 HIV infected patients both on ART and naïve patients

Inclusion criteria

Patients diagnosed to have HIV infection/AIDS after ELISA test being positive.

Exclusion criteria

- 1) Congenital heart diseases
- 2) Pre existing valvular heart disease
- 3) Hypertension

*Correspondence

Dr. Madhu G

Post Graduate , Department of Medicine , BMCRI , Bangalore - 560002, Karnataka , India

E-mail: madhuachar.1993@gmail.com

4) Diabetes mellitus

Investigation required are

- 1) Routine investigations
- 2) Blood HIV 1 and 2 (ELISA)
- 3) CD4, CD8 count
- 4) ECG
- 5) Chest X-ray
- 6) Echocardiography

Sample procedure: 200 HIV infected patients presenting in outpatient department and as inpatients are included in this study. Information is collected and detailed history is taken using preformed proforma.

At the time of admission or follow up, treatment for specific opportunistic infection is initiated and steps are taken to send for all the investigations and detailed clinical examination of the patient is done, focusing more on cardiovascular system.

As cardiovascular manifestations can be symptomatic or asymptomatic cardiovascular work up is done in all patients. Cardiac

manifestations are determined based on clinical, radiological, ECG and Echocardiographic criteria later to correlate the observed study with cd4+ cell counts and whether the patients are on ART or not.

Statistical methods: Student T test was used to match the HIV positive patients on ART and not on ART on the basis of age and sex. Chi-square and Fisher Exact test have been used to test the association of cardiac dysfunction in relation to ART and categories of CD4+ Count. Mann Whitney U, Student T test were used to compare the median CD4+ counts and duration of disease among those with cardiac dysfunction and normal heart

Results and observation

A cross sectional study including 200 ELISA positive HIV infected individuals was taken for a period of two years. Following results and observations were made during the study.

Group I were HIV naïve patients i.e, patients NOT ON ART and Group II patients were NOT ON ART.

Table 1: Age distribution of patients studied

Age in years	Group I	Group II	Total
<20	0(0%)	2(2%)	2(1%)
20-30	6(6%)	9(9%)	15(7.5%)
31-40	35(35%)	38(38%)	73(36.5%)
41-50	43(43%)	38(38%)	81(40.5%)
>50	16(16%)	13(13%)	29(29%)
Total	100(100%)	100(100%)	200(100%)
Mean ± SD	43.08±7.61	42.01±10.05	42.55±8.91

Samples are age matched with P=0.397, student t test

43% of the patients in Group I is in the age group of 41-50yrs, and 38% in the age group of 31-50yrs in group II. On comparing the proportions across the two groups using student T test, it was found the 2 groups were matched by age as the differences were not statistically significant.

Table 2: Gender distribution of patients studied

Gender	Group I	Group II	Total
Female	24(24%)	22(22%)	46(23%)
Male	76(76%)	78(78%)	154(77%)
Total	100(100%)	100(100%)	200(100%)

Samples are gender matched with P=0.737, Chi-Square test

Most of our patients are males in both Group I and Group II, 76% and 78% respectively. No significant difference was noted in the gender distribution.

Table 3: Duration of HIV distribution in two groups of patients studied

Disease Duration	Group I	Group II	Total
<6months	9(9%)	5(5%)	14(7%)
6-12months	4(4%)	4(4%)	8(4%)
1-5 yrs	23(23%)	56(56%)	79(39.5%)
6-10 yrs	2(2%)	26(26%)	28(14%)
>10 yrs	0(0%)	7(7%)	7(3.5%)
Newly diagnosed RVD	62(62%)	2(2%)	64(32%)
Total	100(100%)	100(100%)	200(100%)

56% of the patients in Group II had HIV duration between 1-5 years and most of the patients in Group I are newly diagnosed (62%) and few patients (23%) had disease duration between 1-5 years but who were lost to follow up or not on medications.

Table 4: Types of ART distribution

Type of ART			Total	p value
	Normal Cardiac function	Cardiac dysfunction		
First Line ART	62(89.8%)	26(83%)	88	0.507
Second Line ART	7(10.1%)	5(16.1%)	12	
Total	69	31	100	

*Fishers Exact test used

Among 69 patients with normal cardiac function 62 (89.8%) were on first line ART and among 31 patients with cardiac dysfunction 26(83%) were on first line and 5 (16.1%) were on second line ART.

Table 5: WHO Stage distribution in two groups of patients studied

WHO Stage	Group I	Group II	Total
I	7(7%)	11(11%)	16(8%)
II	38(38%)	45(45%)	83(41.5%)
III	39(39%)	30(30%)	69(34.5%)
IV	16(16%)	14(14%)	30(15%)
Total	100(100%)	100(100%)	200(100%)

41.5% of patients were in WHO stage II of which 45% were in Group II. 39% of the patients belonged to stage III.




Fig 1 a: CD4+ count distribution in each group




Fig 1 b: CD4+ count distribution in each group

The maximum proportion of patients were in the CD4+ Category 51-200 which had 64% and 54% in Group I(NOT ON ART) and II(ON ART) respectively.

Table 6: CD4+ counts distribution in normal and cardiac dysfunction

Attribute	Cardiac function				P value*
	Normal		Cardiac dysfunction		
	Median	Interquartile Range	Median	Interquartile Range	
CD4+ counts	123	77 – 236	102	78 - 278	0.776
Duration Of disease	1.00	0 – 4	0.67	0 - 3	0.140

*Mann - Whitney U test applied due to the skewed distribution of data

CD4+ Counts in patients with normal cardiac function had median value of 123(77-236) and those who had cardiac dysfunction had median value of 102(78-278) but this difference was not found to be statistically significant.

Table 7: Lipid Profile- Comparison in two groups of patients studied

Lipid Profile	Group I	Group II	Total	P value
Total Cholesterol(mg/dl)	102.84±18.07	126.55±45.21	114.70±36.34	<0.001**
LDL (mg/dl)	79.93±21.02	79.63±36.36	79.78±29.62	0.943
VLDL (mg/dl)	46.99±24.62	36.92±21.51	41.96±23.61	0.002**
TGL (mg/dl)	87.07±21.86	121.85±77.02	104.46±59.1	<0.001**
HDL (mg/dl)	22.10±5.31	27.66±10.40	24.88±8.70	<0.001**

We found a statistically significant association between the two groups with total cholesterol,VLDL TGL and HDL.

Table 8: ECG – distribution in two groups of patients studied

ECG	Group I(n=100)	Group II(n=100)	Total(n=200)
Normal	80(80%)	78(78%)	158(79%)
Abnormal	20(20%)	22(22%)	42(21%)
• Sinus tachycardia	8(8%)	10(10%)	18(9%)
• Low voltage complexes	5(5%)	0(0%)	5(2.5%)
• IHD	1(1%)	3(3%)	4(2%)
• LVH	3(3%)	0(0%)	3(1.5%)
• RBBB	1(1%)	2(2%)	3(1.5%)

Majority of our patients had normal ECG (79%), abnormal ECG was found slightly high in Group II patients where the common being sinus tachycardia (10%).




Fig 2 a: WHO Staging and HIV naïve and On ART patients association between Normal and cardiac dysfunction




Fig 2 b: WHO Staging and HIV naive and On ART patients association between Normal and cardiac dysfunction
 44 patients (37.6%) had normal cardiac function who belonged to WHO stage II and 39 patients (47%) had cardiac dysfunction under the same stage but no significant difference was found in the distribution. 62.7% of the patients with cardiac dysfunction were HIV naive and 37.3% were on ART. And statistically significant difference observed between 2 groups (p value <0.003)




Fig 3: Correlation of patients not on ART and on ART with cardiac dysfunction

21 out of 52 patients who were NOT ON ART and 8 out of 31 who were ON ART had pericardial effusion but no significance was elicited (P value 0.17). But 8 patients who were NOT ON ART and 11 patients on ART had TR and also it implies that ART was significantly associated with the occurrence of TR (p value 0.03). Similarly out of 3 patients who were on ART had IHD and it appeared to have significantly associated with ART (p value 0.04).




Fig 4: Correlation of CD4+ counts with cardiac dysfunction




Fig 5: Correlation of CD4+ Counts with cardiac dysfunction in patients NOT ON ART

IHD is absent in all these categories

Among the patients who were NOT ON ART, 85.71% of patients who had pericardial effusion had CD4+ counts till 200. This difference in distribution could have been possibly significant if a larger sample size was chosen. However, 66.67% of patients who had systolic dysfunction had CD4+ count of 200 or less and this association of CD4+ category with SD type of cardiac dysfunction was found to be significant (P 0.003). Though other types of cardiac dysfunction also showed that proportion of patients with lesser CD4+ counts had more cardiac dysfunction, it was not statistically significant.

Discussion

HIV infection has become a chronic and manageable disease. Patients with HIV infection share many cardiovascular risk factors with the general population, but in addition the HIV virus itself, chronic inflammation, opportunistic infections and exposure to ART may increase the cardiovascular risk. Cardiac dysfunction is clinically quiescent in the early stages and later it may become direct cause of death. The present study was carried out at Victoria Hospital and Bowring & Lady Curzon Hospitals attached to Bangalore Medical College to study the cardiovascular manifestations in treatment naïve HIV patients and on treatment (ART) and its

correlation with CD4+ counts.200 patients were considered for the study where 100 patients were NOT ON ART (treatment naïve; Group I) and 100 were on ART (Group II). Mean age group in Group I was 43.08±7.61 years and 42.01±10.05 years in Group II. Most of our patients were males in both the group I and II (76% & 78% respectively) male to female ratio was around 3:1 in both the Groups. Maximum cases were in the age group of 31-50 in both the Groups. In a study by Mehul Merwadi[6] et al., maximum cases were in the age group of 31-40 years of age and male to female ratio was 3:1. Commonest symptoms noted in our study were easy fatigability (53%), weight loss(41.5%) followed by breathlessness (29.5%), cough (28%) and fever(27.5%), but in the study done by Basavaraj[7] et al, noticed commonest symptoms in their study as

fever (82.5%), cough (67.5%) and breathlessness (45%).64% of patients in Group I and 54% of patients in Group II had CD4+ counts in the category 51-200/cumm which is similar to study done by P. Shravan Kumar[8] et al., where maximum no. of patients had CD4+ Count below 200/cumm though no categorization was done below 50/cumm.

Maximum number of patients were in WHO stage III in Group I and Stage II in Group II

Cardiovascular dysfunction in HIV

Cardiac dysfunction was present in 62.7% in patients who are treatment naïve (Group I) and 37.3% were on ART(Group II). Total 83(41.5%) patients of 200 having patients both NOT ON ART and on ART had cardiac dysfunction.

Table 9: showing percentage of patients having cardiac dysfunction in different studies.

	Present study	Mehul Marwadi[6] et al	Kumar SKK[4]et al	Ayaskantha Singh[9] et al
Cardiac Dysfunction	41.5%	49%	58.3%	55.7%

ECG manifestation in HIV infected individuals

Table 10:depicting different ECG findings in different studies.

	Present study	Mehul Marwadi[6]et al.,	Shravan Kumar[8]et al.,	Kumar SKK[4]et al.,
Sinus tachycardia	9%	24%	16%	1.7%
Low voltagecomplexes	2.5%	6%	0.5%	0
IHD	2%	0	3%	1.5%
LVH	1.5%	8%	0.5%	0.3%

X RAY abnormalities in HIV infected individuals

Table 11:Comparing different X ray findings in different studies.

	Present study	Mehul Marwadi[6] et al.,
Cardiomegaly	6%	8%
Right Pleural effusion	7%	6%
B/L reticulonodular pattern	4.5%	12%

Right Pleural effusion was found in 7% of cases followed by Cardiomegaly found in 6% of cases but this results were not similar to other study.

ECHO abnormalities in HIV infected patients

Most common ECHO abnormality found in the present study was Pericardial effusion(34.9%) and pulmonary artery hypertension(34.9%) followed by diastolic dysfunction(28.9%)

Table 12: showing different cardiac dysfunction in different studies

	Present study	Ayaskantha Singh[9]et al.,	Kumar SKK[4] et al.,	Shravan kumar P K[8]et al.,
Pericardial effusion	14.5%	17.4%	13.3%	12%
Dilated cardiomyopathy	5.5%	8.5%	15%	5.5%
Systolic dysfunction	5.5%	22.8%	-	9%
Diastolic dysfunction	12%	8.4%	25%	33%
Pulmonary hypertension	14.5%	11.42%	10%	8%
Mitral regurgitation	8.5%	10%	-	3.5%
Tricuspid regurgitation	9.5%	-	-	-
Ischemic heart disease	1.5%	1.4%	-	-

All the above studies showed cardiac manifestations in patients with HIV but did not specify whether patients were on ART or NOT ON ART.In our study, out of 29 (14.5%) patients having pericardial effusion 21 (72.4%) were NOT ON ART and 8(27.6%) were ON ART but no significant association was found. In a study by Lind A[10] et al., only 0.25% had pericardial effusion who were on ART.We did not find any significant association of ART with Dilated Cardimyopathy in our study. However, one study by Domanski MJ[11] et al., stated that cardiomyopathy would develop was 8.4 times greater in children who had previously used AZT than in those who had never taken AZT.We found that IHD and TR was more common in patients who were ON ART than NOT ON ART (p value 0.049), further studies are warranted to know the causal

association of ART and MI.All the 3 patients who had IHD were on TLE regimen. This result is consistent with a study done by Jens et al., where they've found that the incidence of myocardial infarction increased with increasing exposure to combination antiretroviral therapy (p<0.001).When we tested for association of CD4+ counts with the type of cardiac dysfunction, all showed increased prevalence with decrease in CD4+ counts but only Systolic dysfunction (p 0.048) and IHD (p 0.019) showed statistical significance. Ayaskantha Singh [9] et al., found that patients with CD4 count less than 200/microL had a high prevalence of echocardiographic abnormalities than those with CD4 count more than 200/microL. CD4 count had a significant positive correlation with reduction in EF and Fractional shortening (p<0.001and p<0.002,respectively).Left ventricular

concentric hypertrophy had an inverse relation with CD4+ Count and it was statistically significant in a study by Kumar SKK[4] et al.. Lower CD4+ counts had a higher prevalence of cardiac dysfunction and this association was found to be statistically significant for systolic dysfunction. Pericardial effusion correlation could be significant in a larger sample size. There was no significance in the group of patients ON ART with the CD4+ counts. In a study by, Bijl M et al. 96 patients out of 105 were on ART and 3 of them had Systolic dysfunction with low CD4+ counts. No previous studies have showed groups which have seen the pattern of CD4+ Counts and cardiac dysfunction stratifying equally based on ART and NOT ON ART. In the present study, larger proportion of patients with cardiac dysfunction belonged to WHO stage II and III but no significant association was derived, even in subgroup analysis of Group I and Group II, patients did not show any significant association with cardiac dysfunction and WHO stages. In study by Ayaskantha Singh et al., patients belonged to WHO stage IV more commonly compared to other stages.

Strength of our study

There are very few studies which have divided the sample size equally into those who are treatment naïve and on ART. We have correlated the cardiac dysfunction with WHO clinical staging

Limitations

As the sample size was small, cardiac dysfunction like pericardial effusion did not show statistical significance

Due to resource limitations further evaluation of IHD patients could not be done.

Conclusion

There was significant association of ART on the occurrence of cardiac dysfunction i.e, being on ART or not on ART has significant association with cardiac dysfunction.

2D ECHO is an important diagnostic tool for the evaluation of cardiac dysfunction in these patients.

Cardiac dysfunction is noted with low CD4+Counts and was found to be statistically significant.

Pericardial effusion, diastolic dysfunction, pulmonary arterial hypertension and dilated cardiomyopathy was frequently found in groups NOT ON ART but was not statistically significant.

It is therefore important to undergo 2DECHO at diagnosis and at regular intervals in all HIV infected patients including both ON ART and NOT ART. In the present study sinus tachycardia was the most

common ECG manifestation which also was more commonly found in the other studies mentioned.

References

1. Global Health Observatory (GHO) data. WHO/ UNAIDS Fact Sheet 2016.
2. National AIDS Control Organization (NACO), HIV Facts & Figures 2015.
3. Corallo S, Mutinelli MR, Moroni M, Lazzarin A, Celano V, Repossini A, et al. Echocardiography Detects - Myocardial Damage in AIDS. *European Heart J.* 1998;9(8):887-92.
4. Kumar SKK et al. A study of clinical profile of cardiac dysfunction in patients with HIV infection *Int J Res Med Sci.* 2016;4(12):5149-5153
5. Fauci AS, Lane HC. HIV Disease: AIDS and Related Disorders. *Harrison's Principles of Internal Medicine*, 18th Edition, McGraw Hill Company Inc., New York/Chicago/New Delhi. 2011:1506-87.
6. Marwadi. M, Nileshkumar, Gheewala. G, Rana. J, Bavarva. N. Cardiac manifestations in HIV patients and its correlation with CD+4 cell counts. *National Journal of Medical Research.* 2014;4(3): 244-248.
7. Basavaraj Anita et al, cardiac dysfunction associated with HIV infection. *JAPI.* 2003; 51:1182.
8. Shravan Kumar. P, Siddeswari. R, Sridhar. D, Shakuntala. P, Sitaram et al, Prevalence of Cardiac Manifestations in HIV Infected Patients Correlating with CD4 Count. *International Journal of Scientific and Research Publications.* 2015;4(5):9
9. Singh Ayaskantha, Das Sidhartha, Kumar Dalai Rabindra, Department of Medicine, SCB Medical College and Hospital, Cuttack, India. Study of Cardiac Manifestations in Patients with HIV Infection and Their Correlation with CD4 Count in Indian Population. *International Journal of Clinical Medicine.* 2012; 3: 178-183
10. Lind A, Reinsch N, Neuhaus K, et al. Pericardial effusion of HIV- infected patients - results of a prospective multicenter cohort study in the era of antiretroviral therapy. *European Journal of Medical Research* 2011;16(11):480-483.
11. Domanski MJ, Sloas MM, Follmann DA et al, Effect of zidovudine and didanosine treatment on heart function in children infected with human immunodeficiency virus. *J Pediatr.* 1995;1:2

Conflict of Interest: Nil

Source of support: Nil