

A study to evaluate Lipid profile between type 2 Diabetes mellitus without any complications and type 2 Diabetes mellitus with nephropathy

Md Abu Nasar¹, Ram Ranjan Singh², Brajesh Kumar Singh³, Khursheed Alam^{4*}

¹Associate Professor, Department of Biochemistry, Government Medical College, Bettiah, Bihar, India

²Professor, Department of Biochemistry, Lord Buddha Koshi Medical College, Saharsa, Bihar, India

³Assistant Professor, Department of Medicine, Lord Buddha Koshi Medical College, Saharsa, Bihar, India

⁴Assistant Professor, Department of Surgery, Government Medical College, Bettiah, Bihar, India

Received: 22-08-2021 / Revised: 20-09-2021 / Accepted: 14-10-2021

Abstract

Introduction: Abnormalities of lipoprotein metabolism is associated with Diabetic nephropathy (DN), which can be influenced by derangement of renal function and the degree of metabolic control in diabetes mellitus. **AIM:** Our Aim is to compare the levels of lipid profile, urine microalbumin and glycated haemoglobin (HbA1c) between type 2 Diabetes mellitus (T2DM) without any complications, T2DM with nephropathy and normal controls who were age and sex matched. **Materials and Method:** Study group consisted of 75 individuals of whom 25 were T2DM without any complications, 25 were T2DM with nephropathy and 25 were healthy controls. Total cholesterol (TC), Triglycerides (TG), LDL Cholesterol (LDL-c), HDL Cholesterol (HDL-c), HbA1c and urine microalbumin were measured in these subjects. **Results:** The values of TC, TG, LDL-c, HbA1c and urine microalbumin levels were significantly higher in T2DM without any complications and T2DM with nephropathy when compared to controls. HDL-c level was significantly lower in T2DM without any complications and T2DM with nephropathy when compared to controls. **Conclusion:** T2DM and DN are associated with dyslipidemia which is more pronounced in diabetic nephropathy. Therefore, early detection of dyslipidemia in Type 2 diabetics and accordingly therapeutic intervention could control the resulting cardiovascular or renal complications.

Keywords: Type 2 diabetes mellitus, Diabetic nephropathy, Lipid profile, Urine microalbumin.

This is an Open Access article that uses a funding model which does not charge readers or their institutions for access and distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>) and the Budapest Open Access Initiative (<http://www.budapestopenaccessinitiative.org/read>), which permit unrestricted use, distribution, and reproduction in any medium, provided original work is properly credited.

Introduction

Diabetes Mellitus is a disorder associated with chronic hyperglycaemia resulting from complete or partial decrease in insulin secretion or decrease in biological action of insulin or both [1].

A variety of interrelated lipid and lipoprotein abnormalities is known to be associated with T2DM commonly termed as diabetic dyslipidemia. Hypertriglyceridemia, Low HDL-c, small dense LDL particles is commonly associated with T2DM [2,3].

Deficiency of insulin decreases lipolysis and increases the hydrolysis of stored triglyceride, resulting in greater release of non-esterified fatty acids (NEFA), which is delivered to the liver leading to increased hepatic triglyceride production, which will lead to hepatic VLDL production [4].

Action of insulin on the lipolytic enzyme lipoprotein lipase is reduced which will lead to decreased clearance of triglyceride rich lipoproteins, VLDL and chylomicrons, thus contributing to hypertriglyceridemia in T2DM [5].

Insulin resistance increases the catabolism of HDL in the presence of normal levels of Cholesterol Ester Transport Protein (CETP) and Hepatic Lipase, which promotes the movement of cholesterol ester from HDL to Very Low Density Lipoprotein (VLDL), in the presence of higher VLDL-TG level. HDL-c production is also

decreased secondary to impaired catabolism of VLDL-c and impaired lipoprotein lipase activity [6]. The most common pathology for CKD (Chronic kidney disease) is Diabetes Mellitus. Reduction in lipid abnormalities might preserve glomerular filtration rate, decrease proteinuria, thus reducing the rate of micro and macrovascular complications of diabetes mellitus [7]. This study compared the levels of lipid profile, urine microalbumin and HbA1c between T2DM without any complications, T2DM with nephropathy and age and sex matched normal controls.

Materials and Method

The study population consists of seventy five individuals, who were divided into following groups:

Group I: Diagnosed cases of T2DM without any complications (25)

Group II: Diagnosed cases of T2DM with nephropathy (25)

Group III: Normal controls who were age and sex matched (25)

Inclusion Criteria: Known cases of T2DM without any complications and T2DM with nephropathy aged between 40-70 years.

Exclusion Criteria: Subjects with previous history of thyroid disease, patients on medications such as thyroxine, antithyroid drugs, oral contraceptives and glucocorticoids, pregnancy and coexisting hepatic disease.

Ethical considerations: Approval by the institutional ethical committee was obtained. The objectives of the study were explained to all eligible subjects who participated in the study. Informed consent was sought from all participants.

Sample collection: 7 ml of venous blood specimen was collected from the antecubital vein after fasting for 8-12 hours. Out of which, 2 ml of blood was transferred into EDTA vacutainer and 5 ml blood was transferred into a plain vacutainer. 5 ml of urine sample was

*Correspondence

Dr. Khursheed Alam

Assistant Professor, Department of Surgery, Government Medical College, Bettiah, Bihar, India

E-mail: kalamrxl@gmail.com

collected. The blood samples were subjected to centrifugation and the sera was collected. Total cholesterol is measured by enzymatic method CHOD-PAP (Cholesterol Oxidase-Peroxidase 4- aminoantipyrine) in semiautoanalyzer using commercially available kit.

Serum HDL-c is measured by enzymatic method after precipitation with polyanions in a semiautoanalyzer using commercially available kit.

Friedewald Formula was used to calculate serum LDL- c.

$LDL\ cholesterol\ mg/dl = Total\ cholesterol - [HDL\ Cholesterol + Triglycerides/5]$

Serum triglycerides is measured by enzymatic method GPO-ESPAS (Glycerol 3 Phosphate Oxidase- Peroxidase N-Ethyl-N-Sulfopropyl-n-anisidine) using commercially available kit in semiautoanalyzer.

Ion Exchange Resin method was used to measure HbA1c.

Turbidimetric immunoassay method was used to estimate urine microalbumin in a semiautoanalyzer using commercially available kits.

Statistical analysis: Obtained data was analyzed using SPSS software. Quantitative variables were presented in the form of Mean \pm SD. Analysis of variance (ANOVA) was used to do the comparison between the quantitative variables. Statistical significance was considered to be moderate at p value < 0.05 and p value < 0.01 was considered strongly significant.

Results

Serum TC levels in T2DM without any complications was 255.52 ± 19.75 , in T2DM with nephropathy was 258.56 ± 21.20 and in controls was 163.68 ± 17.68 . T2DM without any complications and T2DM with nephropathy had statistically significant higher levels of total Cholesterol when compared to controls. ($p < 0.001$) (Table 1)

Table 1: Comparison of serum total cholesterol (mg/dl) between the study groups

Study Groups	n	Mean (mg/dl)	SD	F	P
Type 2 diabetics without any complications	25	255.52	± 19.75		
Type 2 diabetics with nephropathy	25	258.56	± 21.20	189.28	< 0.001
Controls	25	163.68	± 17.68		

Serum triglycerides levels in T2DM without any complications was 180.84 ± 18.57 , in T2DM with nephropathy was 182.28 ± 14.16 and in controls was 119.72 ± 25.34 . T2DM without complications and

T2DM with nephropathy had statistically significant higher levels of triglycerides when compared to controls ($p < 0.001$) (Table 2)

Table 2: Comparison of serum triglycerides (mg/dl) between study groups

Study Groups	n	Mean (mg/dl)	SD	F	P
Type 2 diabetics without any complications	25	180.84	± 18.57		
Type 2 diabetics with nephropathy	25	182.28	± 14.16	80.56	< 0.001
Controls	25	119.72	± 25.34		

Serum HDL- c levels in T2DM without any complications was 32.04 ± 7.13 , in T2DM with nephropathy was 31.92 ± 5.13 and in controls was 55.04 ± 11.03 . T2DM without any complications and T2DM

with nephropathy had statistically significant lower levels of HDL when compared to controls. ($p < 0.001$) (Table 3)

Table 3: Comparison of serum HDL (mg/dl) between the study groups

Study Groups	n	Mean (mg/dl)	SD	F	P
Type 2 diabetics without any Complications	25	32.04	± 7.13		
Type 2 diabetics with nephropathy	25	31.92	± 5.13	66.87	< 0.001
Controls	25	55.04	± 11.03		

Serum LDL-c levels in T2DM without any complications was 187.43 ± 22.01 , in T2DM with nephropathy was 189.78 ± 22.93 and in controls was 84.67 ± 19.66 . T2DM without any complications and

T2DM with nephropathy had statistically significant higher levels of LDL-c when compared to controls. ($p < 0.001$) (Table 4)

Table 4: Comparison of serum LDL (mg/dl) between the study groups

Study Groups	n	Mean (mg/dl)	SD	F	P
Type 2 diabetics without any Complications	25	187.43	± 22.01		
Type 2 diabetics with nephropathy	25	189.78	± 22.93	193.27	< 0.001
Controls	25	84.67	± 19.66		

Urine microalbumin levels in T2DM without any complications were 17.24 ± 4.56 , in T2DM with nephropathy was 166.12 ± 34.45 and in controls was 14.12 ± 3.79 . T2DM without any complications and

T2DM with nephropathy had statistically significant higher levels of urine microalbumin when compared to controls. ($p < 0.001$) (Table 5)

Table 5: Comparison of urine microalbumin (mg/dl) between the study group

Study Groups	n	Mean (mg/dl)	SD	F	P
Type 2 diabetics without any complications	25	17.24	± 4.56		
Type 2 diabetics with nephropathy	25	166.12	± 34.45	463.23	< 0.001
Controls	25	14.12	± 3.79		

HbA1c levels in T2DM without any complications were 6.92 ± 1.40 , in T2DM with nephropathy was 8.93 ± 2.35 and in controls was 5.45 ± 0.50 . T2DM without any complications and T2DM with

nephropathy had statistically significant higher levels of HbA1c when compared to controls, which was statistically significant ($p < 0.001$) (Table 6).

Table 6: Comparison of glycated haemoglobin (mg/dl) between the study groups

Study Groups	n	Mean (mg/dl)	SD	F	P
Type 2 diabetics without any Complications	25	6.92	± 1.40		
Type 2 diabetics with nephropathy	25	8.93	± 2.35	29.62	< 0.001
Controls	25	5.45	± 0.50		

Discussion

The world wide prevalence of diabetes mellitus is predicted to increase to 5.4% by the year 2025, most of this increase will occur in developing countries[8].

HbA1c is considered to be the gold standard for measurement of glycemic control. It is also a predictor of diabetic complications: T2DM without any complications and T2DM with nephropathy had increased levels of HbA1c when compared with controls. TC, TG and LDL -c levels are increased and HDL - c is decreased in T2DM without any complications and T2DM with nephropathy, when compared to controls in the present study. These observations are according to the findings of Ejuoghanran OSO et al.

Altered lipid profile in T2DM is due to insulin resistance and defective insulin action on lipoprotein metabolism. Increased lipolysis will increase the synthesis of VLDL and triglyceride rich LDL-c. It will also increase triglyceride synthesis and promote quick breakdown of HDL-c.

Diabetic nephropathy is the consequence of diabetes mellitus characterised by excretion of albumin in urine, increased blood pressure, decreased glomerular filtration rate and high risk of cardiovascular disease. Abnormal lipid profile in diabetes mellitus may lead to worsening of the condition and direct the disease to renal impairment. Elevated lipoproteins and lipids may cause glomerular and tubulointerstitial injury thus contributing to the progression of diabetic nephropathy[11-14].

Conclusion

The present study showed that the common lipid abnormalities associated with patients of T2DM without any complications and T2 DN are hypercholesterolemia, hypertriglyceridemia, elevated LDL-c and decreased HDL -c. Diabetic nephropathy is considered an independent risk factor for cardiovascular disease and altered lipid levels will increase the risk. Routine monitoring of lipid profile should be included in the optimal care for diabetic patients. Lifestyle modifications such as appropriate nutrition, exercise and pharmacological interventions for hyperlipidemia should be implemented for diabetics to prevent or delay its renal manifestations.

References

1. Pasupathi P, Bakthavathsalam G, Saravanan G, Sundarmoorthi R. Screening for thyroid dysfunction in the diabetic/ non – diabetic population. *Thyroid science*. 2008; 3(8):1-6.
2. American Diabetes Association. Diagnosis and classification of diabetes mellitus. *Diabetes Care*. 2005; 28:37-42.
3. Zeqollari A, Spahiu K, Vyshka G, Cakerri L. Lipid profile in diabetes mellitus type 2 patients in Albania and the correlation

- with BMI, Hypertension and hepatoseatosis. *Journal of Family Medicine and community health*. 2014; 1(4):1-5.
4. Valabji J, Elkeles RS. Dyslipidemia in type 2 diabetes – Epidemiology and biochemistry. *The British Journal of Diabetes and Vascular disease*. 2003; 3:184-189.
5. Shen GX. Lipid disorders in diabetes mellitus and current management. *Current pharmaceutical analysis*. 2007; 3:17-24.
6. Mooradian AD, Haas MJ, Wehmeier KR, Wong NC. Obesity related changes in high density lipoprotein metabolism. *Obesity (Silver Spring)*. 2008; 16:1152-116.
7. Collins R, Armitage J, Parish S, Sleight P. Peto: MRC/BHF Heart protection study of cholesterol lowering with simvastatin in people with diabetes: a randomized placebo controlled trial. *Lancet*. 2003; 361:2005-2016.
8. Wild S, Roglic G, Green A, Sicree R, King H. Global Prevalence of Diabetes: Estimates for the year 2000 and projections for 2030. *Diabetes Care*. 2004; 27:1047-1053.
9. Jha P, Shrestha S, Majhi S, Sharma S, Baral N. Glycemic status, Lipid profile and proteinuria in Diabetic nephropathy. *J Nepal Med Assoc*. 2010; 49(2):1-4.
10. Ejuoghanran OSO, Chukwu OE, Christopher SL. The effect of diabetic nephropathy on the lipid profile of diabetics in southern Nigeria. *Journal of Medical sciences*. 2011; 11(4):198-202.
11. Trovati M, Cavalot F. Optimization of hypolipidemic and antiplatelet treatment in the diabetic patients with renal disease. *J Am Soc Nephrol*. 2004; 15:12-20.
12. Gross JL, Azevedo MJD, Silveiro SP, Canani LH, Caramori ML, Zelmanovitz T. Diabetic Nephropathy: Diagnosis, Prevention, and Treatment. *Diabetes Care* 2005; 28 Nosadini R, Tonolo G. Blood glucose and lipid control as risk factors in the progression of renal damage in type 2 diabetes. *J Nephrol*. 2003; 16:42-47:1164-1176.
13. Nosadini R, Tonolo G. Blood glucose and lipid control as risk factors in the progression of renal damage in type 2 diabetes. *J Nephrol*. 2003; 16:42-47.
14. Khan FA, Khan MF, Patil SKB, Jameil N. Estimation of serum copper and magnesium levels in diabetic nephropathy patients. *Asian J Biol Life Sci*. 2013; 2:23-26.

Conflict of Interest: Nil

Source of support: Nil