

## Original research article

**Evaluation of Fungal Diseases of Nose and Paranasal Sinuses in a Rural Tertiary Care Hospital.****Harsh Vardhan<sup>1</sup>, Yashashwi man Anand<sup>2</sup>, Tulika Singh<sup>3</sup>, Tribhuwan Kumar<sup>4</sup>**<sup>1</sup>Senior Resident, Department of ENT, Patna Medical College and Hospital Patna, Bihar, India.<sup>2</sup>Senior Resident, Department of ENT, Patna Medical College and Hospital Patna, Bihar, India.<sup>3</sup>Junior Resident, Department of Pathology, IGIMS, Patna, Bihar, India<sup>4</sup>Associate Professor, Department of ENT, Patna Medical College and Hospital Patna, Bihar, India.**Received: 11-08-2020 / Revised: 15-09-2020 / Accepted: 20-10-2020****Abstract****Aim:** Evaluation of Fungal Diseases of nose and paranasal sinuses in a Rural Tertiary Care Hospital.**Materials and Methods:** The prospective and observational study was conducted in the Department of ENT, Patna Medical College and Hospital Patna, Bihar from December 2017 to November 2018. Total 100 patients who had clinical features suggestive of fungal infections of nose and paranasal sinuses were evaluated with standard pro forma-haematological investigations, radiological procedures, immunological procedures, and pathological diagnostics formed part of the armamentarium. Surgical management and follow-up were done. **Results:** Out of 100 patients study, 35% were male and 65% were female. The majority of cases were in age group between 20-40 years. All patients, in our study, have nasal symptoms 100 (100%). They are nasal obstruction, nasal discharge, postnasal discharge, frequent sneezing, reduced sense of smell (hyposmia) or complete loss of smell (anosmia), and nasal bleeding. Ocular symptoms such as proptosis, epiphora, diplopia blurring of vision in our study were 20%. Fungal culture, in our study, showed out of 100 patients, 33% are *Aspergillus flavus*, 14% *Aspergillus fumigatus*, 10% are *Aspergillus niger*, 3% are *Aspergillus terreus* and 40 were negative. Bilateral disease and involvement of ethmoidal sinus were noted in the majority of cases. **Conclusion:** About 100% of our series of 100 cases were histopathologically proven to be allergic *Aspergillus sinusitis*. CT was found to be highly effective for pre-operative evaluation and intraoperative guidance. Nasal polyposis was a concomitant feature in fungal sinusitis.**Key words:** Allergic fungal rhinosinusitis, *Aspergillus*, Endoscopic sinus surgery, Fungal culture, Sinonasal polyposis

This is an Open Access article that uses a fund-ing model which does not charge readers or their institutions for access and distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>) and the Budapest Open Access Initiative (<http://www.budapestopenaccessinitiative.org/read>), which permit unrestricted use, distribution, and reproduction in any medium, provided the original work is properly credited.

**Introduction**

Fungal sinusitis is the inflammation of the lining mucosa of the paranasal sinuses due to fungal infection.<sup>1</sup> It occurs in people with reduced immunity. The maxillary sinus is the most commonly involved. Fungi responsible for fungal sinusitis are *Aspergillus fumigatus* (90%), *Aspergillus flavus*, and *Aspergillus niger*. Fungal sinusitis occurs most commonly in middle-aged populations. Diabetes mellitus is the most common risk factor involved.<sup>2</sup> The most common causes of sinus infections are linked to viral and bacterial infections. However,

a fungal sinus infection differs in that it refers to an infection of any or all of the four paranasal sinuses caused by a fungal growth inside the body. Fungi can be found both in nature, in your home and work environments, and in the air. Sometimes when fungal material or debris is inhaled into the nose and lungs, it may cause an infection.

There are two forms of fungal sinus infections: invasive and non-invasive. Non-invasive infections do not spread anywhere besides the sinuses, and occur in patients who have normal immune systems. However, invasive fungal sinus infections can occur in those patients whose immune systems are not as strong as the average person. An invasive sinus infection can become life-threatening in a very short amount of time, meaning that immediate treatment is critical. The most common site of fungal infection in man is the lungs

---

*\*Correspondence***Dr. Harsh Vardhan**

Senior Resident, Department of ENT, Patna Medical College and Hospital Patna, Bihar, India.

E-mail: [dr.harshvardhan30@yahoo.co.in](mailto:dr.harshvardhan30@yahoo.co.in)

with or without hematogenous spread to other organs. However, the localized fungal infection can also occur in the upper respiratory tract and is more common than was previously suspected.<sup>3-5</sup>

Most fungal species which are pathogenic to human cause opportunistic infection and only dermatophytes are transmissible from host to host. The incidence of infections and death due to fungi has been grossly underestimated moreover the list of fungal species capable of producing disease in immune compromised person is increasing rapidly. In an era with AIDS, broad spectrum of antibiotics, cytotoxic drugs and the organ transplantation, fungal infection which affect the nose and the sinus are candidiasis, rhinosporidiosis. *Aspergillus*, *phycomycosis*, *actinomyces*, *coccidioidomycosis*, *histoplasmosis*, *cryptococcosis*, *blastomycosis*, *sporotrichosis*, and *nocardiosis*. Martin and Berson noted a high incidence in South Africa which they attributed particularly to malnutrition the largest series of case involve.<sup>6-8</sup>

In our study, the fungal infections mainly presented with nasal polyps, nasal block, nasal discharge, headache, and proptosis mimicking benign or malignant tumors of the nose and paranasal sinuses

#### Material and Methods

The prospective and observational study was conducted in the Department of ENT, Patna Medical College and Hospital Patna, Bihar from December 2017 to November 2018.

The study protocol was reviewed by the Ethical Committee of the Hospital and granted ethical clearance.

#### Methodology

A total of 100 patients who had clinical features suggestive of fungal infection of nose and paranasal sinus were evaluated using a standard pro forma and underwent the following investigate procedures systematically as and when needed.

Haematological Investigations complete with hemogram, blood sugar level, serum electrolyte, serum protein, blood grouping, etc., were done as preliminary investigations to assess the general health condition as well as to rule out any underlying disorders.

Relevant X-ray of the nose and paranasal sinuses was taken for all patient and those who were provisionally diagnosed as fungal granulomas were subjected to CT scanning of the nose, paranasal sinuses, and brain with contrast enhancement studies.

The clinical diagnosis of nasal fungal infection, which was confirmed using microbiological tests by the Consultant Microbiologist. Thereafter, the Consultant ENT directed the patient to the principal investigator for administration of tests within the ENT department.

#### Endoscopic Sinus Surgery

Postoperatively patients were advised to come for regular follow-up. The nasal douching was given to every patient for the 5<sup>th</sup> post-operative day after the first post-operative endoscopic examination and cleaning. Patient was treated by beclometasone aqueous nasal spray, antihistamine, and vitamins.

The patients were requested to come for follow-up on the 15<sup>th</sup> post-operative day for endoscopic examination and cleaning and whenever possible thereafter (usually once in a month). The patient with allergic *Aspergillus sinusitis* did not require antifungal therapy. Antifungal therapy was given based on the type of fungal infection and its invasiveness (mucormycosis).

#### Statistical Analysis

The recorded data was compiled and entered in a spreadsheet computer program (Microsoft Excel 2010) and then exported to data editor page of SPSS version 20 (SPSS Inc., Chicago, Illinois, USA). Descriptive statistics included computation of percentages, means and standard deviations were calculated.

#### Results

**Table 1: Age/sex/incidence N=100**

Age	Sex		Incidence		
	Male	Female	Male (%)	Female (%)	Total
Below 20	6	12	6	12	18
20-30	11	23	11	23	34
30-40	15	25	15	25	40
Above 40	3	5	3	5	8
Total	35	65			

**Table 2: Clinical symptoms N=100**

Symptoms	Number of patients	%
1. Nasal Nasal obstruction Nasal discharge Post nasal Discharge	100	100
2. Headache	70	70
3. Ocular Proptosis Epiphora Diplopia Ophthalmoplegia	20	20

**Table 3: Histopathology and fungal culture N=100**

Causative organism	Number of patient	%
<i>Aspergillus flavus</i>	33	33
<i>Aspergillus fumigatus</i>	14	14
<i>Aspergillus niger</i>	10	10
<i>Aspergillus terreus</i>	3	3
No growth	40	40

**Table 4: CT scan of nose and sinus N=100**

Sinus involvement	Number of patients	%
Maxillary sinus	81	81
Ethmoidal sinus	77	77
Frontal sinus	47	47
Sphenoidal sinus	55	55
All sinuses	31	31
Orbital	17	17

**Table 5: Unilateral/bilateral comparison study N=100**

Sides of nose and sinuses	Number of patients	%
Right	14	14
Left	18	18
Unilateral	32	32
Bilateral	68	68

**Table 6: Complications N=100 (By A disease process)**

Complication	Number of patient
Intraoperative hemorrhage	9
Cerebrospinal leak	Nil
Synechia	30
Periorbital ecchymosis	10

**Table 7: Follow-up and recurrence**

Number of cases	Month of follow-up	Number of recurrence
26 cases	18 months	9
52 cases	12 months	3
22 cases	6 months	-

## Discussion

The fungal diseases of the nose and paranasal sinuses encompass not one disease entity but a multitude of an entire spectrum of different diseases. We have studied different disease causes, namely allergic *A. sinusitis* (100 cases). Although the treatment of these diseases is vastly different, the presentation and clinical features are quite similar and thus they could be studied together. We have attempted to study these diseases under the common heading highlighting the important difference whenever required.<sup>9-11</sup>

The majority of cases in our study were between the age group of 20-40 years. They constitute 74% of a total number of cases. This was followed by the age group below 20 years who constitute 18 % total number of cases. This compares favourably with these studies. In our study, there was a clear female preponderance numbering 65 out of 100 cases (65%) and male 35 out of 100 (35%) reported data by Waman *et al.* showed female preponderance with allergic *A. sinusitis* in our study was 65% female ratio . This corresponds well with the previously mentioned study.

All patients in our study had nasal symptoms. The nasal symptoms included nasal obstructions, nasal discharge frequent sneezing, reduced smell (hyposmia) complete loss of smell (anosmia), and nasal bleeding. The next most common symptom was a headache was seen in 70% of our patient. The next most common symptoms were ocular symptoms such as epiphora, diplopia and blurring of vision comprising about 20%. Various other studies showed that the common symptoms in allergic *A. sinusitis* are chronic nasal obstruction and postnasal discharge.<sup>12,13</sup> These findings compare favorably with our studies.

In our study, out of 100 patients, all 100 patients presented with nasal polyps, fungal mass (100%). The ocular sign such as proptosis diplopia and ophthalmoplegia was seen 20%.

A total of 100 different fungal diseases have been reported in fungal sinusitis. *Aspergillus*, ubiquitous fungus of the class ascomycetes is the most commonly encountered fungus in the environment and is the most common species encountered in fungal sinusitis generally and presumably in allergic fungal sinusitis. The latter is largely based on histopathological finding of fungi with morphologic features similar to *Aspergillus* and not on the basis of culture documentation. In our series 100% of fungal sinusitis was histopathologically proven to be aspergillus. In our study shows 100% allergic *A. sinusitis*. Klossek *et al.* in his case series of 100 cases documented that 94% were histopathologically proven allergic *A. sinusitis* various other organisms have reported as pathogens in allergic *A. sinusitis* caused by different fungi. *Bipolaris*

*specifera* B, Australians, *Aspergillus*, *Alternaria* and *Curvularia lunata*. The identification of these fungi may be related to the improved ability of microbiology laboratories to identify the diverse hyphae with variation in the conical pores.<sup>14,15</sup>

In our study, all the cases of allergic *A. sinusitis* were sent for the fungal culture. In all the cases the material sent for culture were fungal mass taken from the infected sinus cavity. out of 100 cases 60 were culture positive for aspergillus and remaining 40 were negative. Out of 60 percent 33% are *Aspergillus flavus*, 14% *Aspergillus fumigates*, 10% are *Aspergillus niger*, 3% are *Aspergillus terreus*. Rhinomucormycosis was no present in our study in histopathological examination.

In our studies, no other species of fungi was identified either on the HPE or cultural examination. All the patients in this series underwent CT scan preoperatively, magnetic resonance imaging (MRI) scan was not considered due to the high cost factor and relatively low amount of extra information in cases of fungal diseases of nose and paranasalsinus.

All patients in our series with allergic *A. sinusitis* demonstrated areas of high alteration centrally within involved sinus by CT. These areas corresponded to surgical findings of thick allergic mucin. Some cases demonstrate a starry sky pattern of material, which appeared to be calcium densities on bone windows. CT scanning has been very useful in defining the full extent to the disease. *A. sinusitis* often has a mixture of high and low-density areas within the sinuses. Bone windows allow a very accurate assessment of possible invasion.

In general, only one series in involved with *Aspergillus* most commonly the maxillary sinus. In our study, maxillary sinus 81 is the common involvement. Next ethmoid 77%, frontal sinus 47%, sphenoid sinus 55%, and all sinuses involvement 31%.<sup>16</sup>

In our study, 100 cases were operated by endoscopic sinus surgery. Endoscopic sinus surgery with less morbidity and mortality, clearance was total and recurrence rate is almost minimal in our steady. Even though in our cases, we had no complication acquired in functional endoscopic sinus surgery. Only in 12 cases recurrence was noted. None of the patients developed complication and patient were discharged next day itself. This correlates well with the previous study.

We treated our patient with steroids both topically and systemically. The use of tropical intranasal steroids id routine, and we restrict the use of systemic steroids. It is our experience that the tropical intranasal steroids alone when taken regularly are effective in preventing recurrence of the disease. However feels that tropical

intranasal steroids and effectively only after a course of oral corticosteroids.

Antifungal agents were not used in any of our cases with allergic *A. sinusitis*. Similar reports have been published by many authors regarding the endoscopic approach is the sole approach in the treatment of allergic *A. sinusitis*. However, some author feels that external approach definitely has its place in the treatment of this condition especially in cases of orbital (or) intracranial extension of this disease.

Complications of endoscopic sinus surgery have been as major and minor according to the degree of morbidity and treatment needed to prevent permanent serious sequelae. Complications seen in our study includes intraoperative hemorrhage in 9 cases (9%) and no cerebrospinal leak. Pneumocephalus and other reported major complication (Markmay *et al.*, 1994) includes orbital hematoma. Loss of vision, diplopia, epiphora, meningitis, brain abscess, and focal brain hemorrhage which were not seen in our study.

Intracranial complication can be prevented by not disturbing the mucosa lying against the roof of the ethmoid sinus. It is also worth remembering that the vertical bony wall of olfactory groove where the middle turbinate attaches to the roof of the ethmoid sinus may be extremely thin and should be avoided. We feel that two other guidelines may help to prevent cerebrospinal fluid leaks.

Instrumentations or suction cannulas should be placed into the nose or sinuses only under endoscopic guidance. The basal lamella should be entered at a point farthest from the roof of the ethmoids posteriorly and inferiorly rather than anteriorly and superiorly.

Intraoperative hemorrhage severe enough to require blood transfusion is rare in our review none of them require blood transfusion. We agree with other reported studies that this kind of preoperative bleeding is mostly from the interruption of the sphenopalatine artery as it courses over the face of the sphenoid sinus, just above the arch of the posterior nasal choanae.

The most frequently encountered minor complication in our study 30% (30 cases) were synechiae. This adhesion was usually seen between the middle turbinate and septum or lateral wall of nose careful handling of the tissue during surgery minimizes the chance of contact between the two adjacent raw surfaces. Careful post-operative cleaning of the sinus cavity will also help in the prevention of adhesion of the 100 patient in our study, 30 had synechiae which were released in the outpatient department, and there was no recurrence.

Periorbital ecchymosis is the next minor complication and a total of 10%. These complications were seen after the endoscopic sinus surgery. This occurs usually due to violation of lamina papyracea. Other reported

studies that the violation of the lamina papyracea occurs most commonly with uncinectomy during endoscopic sinus surgery.

Post-treatment endoscopic surveillance is essential for long-term success since recurrent disease is common. Furthermore, the patient symptoms alone are not a satisfactory measure for persistent/recurrent disease.

In this series, 26 of our patient were followed up for a period of 18-month after surgery. 52 patients were followed up for a period of 12-month and another 22 patients followed up for 6 months. We have not lost any patient during the follow-up treatment and proper medication. 12 recurrences were noted within 18 months.

The complete and radical removal of fungal debris and careful regular follow up with intranasal steroids and if required systemic steroids when employed judiciously will result in the best long-term result after surgery.<sup>17-19</sup>

### Conclusion

About 100% of our series of 100 cases were histopathologically proven to be allergic *Aspergillus sinusitis*. CT was found to be highly effective for pre-operative evaluation and intraoperative guidance. Nasal polyposis was a concomitant feature in fungal sinusitis. The data generated from the present study and the reported literature suggests that combination of medical and surgical line of treatment improved the prognosis significantly. Early detection, proper and adequate dose of antifungal agents, timely surgical intervention improve the survival rate in the disease of sinonasal fungal infections.

### Reference

1. Fungal Sinusitis: Background, History of the Procedure, Problem. eMedicine. Retrieved 25 November, 2016.
2. Karthikeyan P, Nirmal V Coumare. Incidence and presentation of fungal sinusitis in patient diagnosed with chronic rhinosinusitis. Indian Journal of Otolaryngology Head and Neck Surgery. 2010; 62(4):381-5. doi:10.1007/s12070-010-0062-0. PMC 3266098. PMID 22319697
3. Azadeh B, Baghoumian N, el-Bakri OT. Rhinosporidiosis: Immunohistochemical and electron microscopic studies. J Laryngol Otol 1994;108:1048-54.
4. Bader G. Aspergillus sinusitis of dental origin. Rev Odontostomatol (Paris) 1989;18:345-53.
5. Bassiouny A, Maher A, Bucci TJ, Moawad MK, Hendawy DS. Noninvasive antromycosis: (Diagnosis and treatment). J Laryngol Otol 1982;96:215-28.

6. Becker MH, Ngo N, Beranbaum SL. Mycotic infection of the paranasal sinuses. Radiographic manifestations. *Radiology* 1968;90:49-51.
7. Brantdwein M. Histopathology of sinonasal fungal disease. *Otolaryngol Clin North Am* 1993;26:949-81.
8. Chapnik JS, Bach MC. Bacterial and fungal infections of the maxillary sinus. *Otolaryngol Clin North Am* 1976;9:43-54.
9. Davis BD. Fungi in Microbiology. 2nd ed. Ch. 43. Farnham Royal: Commonwealth Agricultural Bureaux; 1973. p. 964.
10. Finn DG, Farmer JR, Durham NC. Chronic mucormycosis. *Laryngoscope* 1982;92:61-763.
11. Ritter FN. The Para Nasal Sinuses Anatomy and Surgical Technique. St. Louis: CV Mosby; 1978.
12. Glass RB, Hertzanu Y, Mendelsohn DB, Posen J. Paranasal sinus aspergillosis. A case report with computed tomogram findings. *J Laryngol Otol* 1984;98:199-205.
13. Green WR, Font RL, Zimmerman LE. Asperilosis of the orbit. Report of ten cases and review of the literature. *Arch Ophthalmol* 1969;82:302-13.
14. Hora JF. Primary aspergillosis of the paranasal sinuses and associated areas. *Laryngoscope* 1965;75:768-73.
15. Iwamoto H, Katsura M, Fujimaki T. Mycosis of the maxillary sinuses. *Laryngoscope* 1972;82:903-9.
16. Jahrsdoerfer RA, Ejercito VS, Johns MM, Cantrell RW, Sydnor JB. Aspergillosis of the nose and paranasal sinuses. *Am J Otolaryngol* 1979;1:6-14.
17. Lavelle WG. Aspergillosis of the sphenoid sinus. *Ear Nose Throat J* 1988;67:266, 268-9.
18. Mahgoub ES. Mycoses of the Sudan. *Trans R Soc Trop Med Hyg* 1977;71:184-8.
19. Savetsky L, Waltner J. Aspergillosis of the maxillary antrum. Report of a case and review of the available literature. *Arch Otolaryngol* 1961;74:695-8.

**Conflict of Interest: Nil**

**Source of support: Nil**