

**Clinical profile and outcome of obstetric referral pattern: an observational study****Priyanka Bharti****Senior Resident, Department of Obstetrics and Gynaecology, Darbhanga Medical College and Hospital, Laheriasarai, Darbhanga, Bihar, India****Received: 22-06-2020 / Revised: 20-7-2020 / Accepted: 10-08-2020****Abstract****Aim:** To observe clinical profile and outcome of obstetric referral pattern.**Materials and methods:** A prospective study was conducted in the Department of Obstetrics and Gynaecology, Darbhanga Medical College and Hospital, Laheriasarai, Darbhanga, Bihar, India from 1 year, Thorough history was taken; complete physical and obstetric examination and relevant investigations were done. Management of the patient, clinical course, mode of delivery, both maternal and perinatal outcomes were documented. **Results:** A total of 160 cases were referred to the higher centre due to various reasons. Maximum number of cases in present study were in the age group of 20-30 years comprising 102(63.75%) of total cases. Majority of the referral cases were primigravida 85 (53.12%). Out of 160 referred cases, 127 (79.37%) delivered 19 (11.87%) were treated conservatively. In 14 patients (8.76%) either abortion occurred or medical termination of pregnancy was done or there was ectopic pregnancy or tears which were managed according to set protocol depending upon the gestational age at diagnosis. **Conclusion:** The present study has shown that improper antenatal and intranatal care at the periphery level is responsible for poor maternal and perinatal outcome. Hypertensive disorders of pregnancy have been one of the commonest causes of referral among high risk obstetric patients which can be better dealt at the tertiary care centre.**Keywords:** Hypertensive disorders

This is an Open Access article that uses a fund-ing model which does not charge readers or their institutions for access and distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>) and the Budapest Open Access Initiative (<http://www.budapestopenaccessinitiative.org/read>), which permit unrestricted use, distribution, and reproduction in any medium, provided the original work is properly credited.

**Introduction**

Women die every year in India<sup>1,2</sup> which contribute 20-25% of all maternal deaths in the world<sup>3</sup>. One estimate shows that with one maternal death, 15% pregnancies develop complication which necessitates tertiary obstetric care<sup>4</sup> and the vast majority of maternal deaths and injuries are avoidable when women have access to health care before, during and after childbirth. Of course there is improvement in maternal and child healthcare after the millennium declaration 2000, but there are lacuna across different states, Kerala being the most outstanding and Uttar Pradesh the worst performer.<sup>4,5</sup>

Emergency obstetric transfers should be carried out effectively and efficiently to avoid maternal and foetal morbidity and mortality.

**\*Correspondence****Dr. Priyanka Bharti**

Senior Resident, Department of Obstetrics and Gynaecology, Darbhanga Medical College and Hospital, Laheriasarai, Darbhanga, Bihar, India

E-mail: [Priyanka.sarraf@outlook.com](mailto:Priyanka.sarraf@outlook.com)

An institution referral is when a pregnant woman seeks care at a lower level health facility (basic emergency obstetric care) and is referred onwards to a higher level health facility (comprehensive emergency obstetric care). Referral systems have been considered to be an important component of health systems in developing countries since the emergence of primary healthcare. Referral is especially important within obstetrics due to the high numbers of professionals who support a woman through pregnancy and birth, the speed with which action often needs to be taken and the global burden of maternal mortality.<sup>6</sup> The World Health Organization estimates that at least 88–98% of maternal deaths can be averted with timely access to existing, emergency obstetric care using effective and efficient referral systems. A good and well sustained referral system needs referral- protocols, improved support (especially transportation), community awareness, and feedback system.<sup>7</sup>

Due to lack of awareness and absence of regular antenatal care, the critically ill patients are referred late and sometimes in moribund conditions with multiple

organ damage. Timeliness and appropriateness of referral is an important factor in the ultimate outcome of the patients<sup>8</sup>. Referral services for identification and referral of high risk pregnancies are an integral part of maternal and child health services. For a large majority of developing countries this aspect of health system remains weak.<sup>9</sup> Although most obstetric complications (defined as acute conditions such as postpartum haemorrhage, sepsis, eclampsia, and obstructed labor that can cause maternal death cannot be predicted, the majority can be treated with timely provision of a package of evidence-based interventions known as emergency obstetric care (EmOC)<sup>10-12</sup>. The availability of EmOC is considered to be an indicator of how well a health system is prepared to manage conditions leading to acute maternal morbidity and mortality.<sup>13,14</sup> Em OC refers to elements of obstetric care needed for management of complications during pregnancy, delivery and postpartum period, skilled personnel, equipment and support services. EmOC services are of paramount importance in reducing maternal mortality and morbidity.<sup>15</sup> It is still recommended to electively refer pregnant woman with previous caesarean section, breech presentation, transverse lie, multiple gestation, hypertension and severe anaemia for delivery before any complication arise to a health care centre where all the facilities to deal with the complications are available.<sup>15</sup> With this background present study was undertaken to evaluate the maternal and perinatal outcome in referred cases.

### Material and Methods

A prospective study was conducted in the Department of Obstetrics and Gynaecology, Darbhanga Medical College and Hospital, Laheriasarai, Darbhanga, Bihar, India.

The study protocol was reviewed by the Ethical Committee.

### Methodology

Using a pre structured designed questionnaire, socio demographic details, medical co-morbidities, indications for referral maternal and foetal were obtained. Referral slips were analysed and source of referral, distance travelled and mode of transport and referral-arrival interval, documentation patterns were sought. Patient referred while in labour were noted. Gestational age at referral and mode of delivery was highlighted. Intra partum variables and surgical morbidities were evaluated. To know perinatal outcome, APGAR score was noted, if needed NICU admission cause for it was noted.

### Statistical Analysis

The recorded data was compiled and entered in a spreadsheet computer program (Microsoft Excel 2010) and then exported to data editor page of SPSS version 19 (SPSS Inc., Chicago, Illinois, USA). Descriptive statistics included computation of percentages.

### Results

**Table 1: Age Distribution**

Age years	N=160	%
Below 20	3	1.87
20-30	102	63.75
30-40	44	27.5
Above 40	11	6.88

**Table 2: Parity-Wise Distribution**

Parity	N=160	%
Primigravida	85	53.12
Multigravida	66	41.25
Grand multigravida	9	5.63

**Table 3: Outcome of the Antenatal Cases**

Outcome of ANC	No. of cases	%
Delivered	127	79.37
Abortion /ectopic	14	8.76
Conservative	19	11.87
Total	165	100

**Table 4: Mode of Delivery.**

Mode of Delivery	N=127	%
Normal Delivery	81	63.78
LSCS	46	36.22

**Table 5: Reasons for NICU Admission.**

Reason for admission	No. of cases=30	%
Preterm care	13	43.33
Meconium aspiration syndrome	8	26.67
Jaundice	3	10
Sepsis	2	6.67
Transient tachypnoea of newborn	1	3.34
Low birth weight	1	3.33
Asphyxia+ death	1	3.33
Hypoglycaemia	1	3.33

**Table 6: Causes of referral**

Parameter	N=160
Preterm Labour	13
PROM	26
Pre-eclampsia and related condition	22
Cardiac disease	2
Crossed dates	10
Prev LSCS	7
Antepartum hemorrhage	9
Postpartum hemorrhage	8
MSAF	16
Malpresentation	3
Non availability of blood	7
Non availability of doctor	5
No details	7
Ectopic	2
Short stature	3
Anaemia	10
Fetal distress	10

**Table 7: Most common period during which patients were referred**

Period of pregnancy	No. of cases=160	%
Intrapartum	106	66.25
Antenatal	37	23.13
Postpartum	17	10.62

## Discussion

Labour is a physiological process, but it carries an inherent risk of complications. Obstetrical care in the western world is at its peak. But in developing countries it is still at docks due to illiteracy, male dominant society and untrained birth attendants. Majority of the population living in the rural areas do not have accessibility to the maternity centres and may develop life threatening complications during labour.<sup>16</sup> The death of a woman in childbirth is a tragedy, an unnecessary and wasteful event that carries with it the huge burden of grief and pain. Pregnancy is not a disease and pregnancy related morbidity and mortality are almost preventable. Identification of atrisk patients and obstetrics emergencies and timely referral is of immense importance.<sup>15</sup> Morsheda Banu et al on assessing the overall age distribution found that the majority (74%) of the respondents were between 20-35 years.<sup>17</sup> In the study conducted by Prakriti Goswami et al, the maximum number of patients were in the age group of 20-30 years (78%).<sup>15</sup> Similar kind of results was observed in the present study. Gupta PR et al found 52.17% patients were primigravida<sup>18</sup>, Prakriti Goswami et al found 47% patients were primigravida<sup>15</sup>, Morsheda Banu et al had found that 50% of women were primigravida<sup>17</sup>, which is comparable to the 53.12% primigravida cases found in the present study. 66.25%, 23.13%, 10.62% of the cases were referred to our hospital in their intapartum, antepartum and postpartum period respectively. Similar results are found by Prakriti Goswami et al where 56%, 30% and 14% of cases were referred in intrapartum, antepartum, postpartum period respectively<sup>15</sup> and also by Devinneni K et al in their "Study of spectrum of referral pattern at a tertiary teaching hospital toward better obstetric care".<sup>19</sup> Out of the 127 cases who delivered at our institute majority of the babies were delivered 63.78% delivered normally, while 36.22% underwent caesarean section. Similar results were obtained by Gupta PR et al where 69.48% of the patients underwent vaginal delivery and 22.75% of the patients had LSCS.<sup>18</sup> The caesarean section rates in the present study was found to be similar to the study conducted by Goswami et al (28%).<sup>15</sup> In the present study, it can be concluded that the rate of caesarean section is substantially high in referred cases. Patel HC et al in their study found that causes of referral were pre-eclampsia (16%) and meconiumstained liquor (5%)<sup>20</sup>. Sabale et al in their study found that preeclampsia and related conditions were a major indication for referral (25.79%).<sup>21</sup> Rathi Charu et al noted that a majority of the cases were referred for preeclampsia and related conditions (26%), preterm labour (26%) and medical disorders

complicating pregnancy (21%).<sup>22</sup> In the present study, premature rupture of membranes was the most common cause of referral (16.25%). This is followed by pre-eclampsia and related conditions (13.75%), and meconium stained liquor (10%). Previous caesarean sections were the cause of referral in 4.38% of cases in the present study which is similar to the study conducted by Goswami P et al(6%)<sup>15</sup>, Khatoun A et al (15%)<sup>23</sup> and Gupta PR et al(7.62%).<sup>18</sup> The patients with previous caesarean section are referred to higher centres from PHC/CHC due the unavailability of operation theatre, gynaecologist, anaesthesiologists, trained staff or basic infrastructure deficit.<sup>15</sup> In the present study, 7.5% of cases were referred due to non availability of blood and doctors; this can be compared to the study conducted by Goswami P et al. where 16.87% of the cases were referred for the same reason. Government should take measures to improve health infrastructure facilities, make provisions for developing new blood banks and appoint trained gynaecologists in the peripheries to reduce the burden on tertiary centres.<sup>15</sup> 11.87% of the total referred cases were managed conservatively and discharged. This rate is similar to the study conducted by Gupta PR et al (7.76%)<sup>18</sup>, Poornima M et al (11%)<sup>24</sup>, Goswami P et al (24%).<sup>15</sup> Here arises the concept of day care management of referral cases at tertiary care institute which might be helpful in reduction of burden of tertiary care institute.<sup>18</sup> Khatoun A et al had in their study reported 87% live births, 13% still births, 26.5% preterm births.<sup>23</sup> Rathi Charu et al in her study found that 90% were live births and 9% were still births.<sup>22</sup> Poornima M et al in her study reported 91% live births and 9% still births.<sup>24</sup> In the present study, 99.38% were live births. In the present study 3.33% had birth asphyxia and 18.75% had NICU admissions, which is comparable to the study conducted by Poornima M et al where 27% of the babies needed NICU admissions.<sup>24</sup> In the present study 43.33% of the NICU admissions were for preterm care, 26.67 for meconium aspiration syndrome, followed by 10% for neonatal jaundice. These results are similar to the findings found in the study conducted by Poornima M et al where 47% of NICU admissions were for preterm care, 28% for respiratory distress.<sup>24</sup> The high rate of NICU admission is due to preterm delivery. The neonatal death rate is 3.33% which is similar to the neonatal death rate in study conducted by Poornima M et al (8%)<sup>24</sup>, Sabale U et al (10.23%)<sup>21</sup>, Gupta PR et al (4.43%).<sup>18</sup> In contrast to this, the study conducted by Rathi Charu et al had a neonatal mortality rate of 28.23%.<sup>22</sup>

### Conclusion

Childbirth is a normal physiologic process, but emergencies can arise anytime. The present study has shown that improper antenatal and intranatal care at the periphery level is responsible for poor maternal and perinatal outcome. Hypertensive disorders of pregnancy have been one of the commonest causes of referral among high risk obstetric patients which can be better dealt at the tertiary care centre. Health care workers should be provided with the checklist; also administration of a dose of magnesium sulphate must be done in all cases of eclampsia and severe pre-eclampsia prior to referral. The quality of information provided on the referral slips leaves a lot to be desired. Many of the slips had no vital data like blood pressure, pulse rate, temperature, and presence of foetal heart sounds. Also information regarding treatment given before referral of the case was missing in most of the referral slips. The cause of referral was unknown in 3% of cases. This call for a more stringent documentation in referral slips and more coordination between the referral units and the higher centres are required.

This study also recommends the development of a standard referral protocol, proper training in this regard and universal adherence to this in practice. Health education and awareness by mass media and non-government organisations can improve the health and social status of women in rural areas.

### Reference

1. National Family Health Survey (NFHS-2) 1998-99. Vol. 196. Bombay: International Institute of Population Services; 2000. pp. 247-51.
2. National Family Health Survey (NFHS-3) 2005-2006. Vol. 196. Bombay: International Institute of Population Services; 2006. pp. 247-51.
3. Pal A, Ray P, Hazra S, Mondal TK. Review of changing trends in maternal mortality in a rural medical college in West Bengal. *J Obstet Gynecol India*. 2005;55:521-4.
4. Mohapatra A, Gomare M. A critical appraisal of the maternal and child health scenario in a metropolitan city in India with reference to achievements of millennium development goals. *J Family Med Prim Care*. 2019;8:995-1001.
5. Selected socioeconomic statistics of India 2017.
6. World Health Organisation, UNICEF, UNFPA, World Bank, UN Population Division. Trends in maternal mortality: 1990 to 2013. Estimates by WHO, UNICEF, UNFPA, the World Bank and the United Nations Population Division. Geneva: World Health Organisation; 2014.
7. Stefanini A. District hospitals and strengthening referral systems in developing countries. *World hospitals and health services: World Hosp Health Serv*. 1994;30:14-9
8. Gupta PR, Chaudhari SN, Gonnade NV. Maternal and fetal outcome in referred patients to tertiary care centre, *Sch J App Med Sci*. 2016;4(5C): 1624-31
9. Park K Textbook of Social and Preventive Medicine 17th ed. Jabalpur. Banarasidas Bhanot Publishers 2000:632
10. Koblinsky M, Chowdhury ME, Moran A, Ronsmans C. Maternal morbidity and disability and their consequences: neglected agenda in maternal health. *J Health Popul Nutr*. 2012;30 (2): 124-30.
11. Lee AC, Lawn JE, Cousens S, Kumar V, Osrin D, Bhutta ZA. Linking families and facilities for care at birth: what works to avert intrapartum-related deaths? *Int J Gynaecol Obstet*. 2009;107(1):S65-85.
12. World Health Organisation. Monitoring the Building Blocks of Health Systems: A Handbook of Indicators and their Measurement Strategies. Geneva: WHO; 2010
13. Travis P, Bennett S, Haines A, Pang T, Bhutta Z, Hyder AA. Overcoming health-systems constraints to achieve the Millennium Development Goals. *Lancet*. 2004; 364 (9437): 900-6.
14. Paxton A, Bailey P, Lobis S. The United Nations Process Indicators for emergency obstetric care: Reflections based on a decade of experience. *Int J Gynaecol Obstet*. 2006;95(2):192-208.
15. Goswami P, Bindal J, Chug N. To study pattern of obstetric cases referred at tertiary care centre in Central India. *Int J Reproduct Contracep, Obstet Gynecol*. 2017; 6(6): 2370-2374.
16. Sumera S, Nuzhat R, Sadia A. Fetomaternal outcome in cases referred to tertiary care hospital after trial of labour. *JSZMC*. 4(2):444-447.
17. Morsheda B, Shamsun N, Hashima EN; Assessing the MANOSHI Referral System Addressing Delays in Seeking Emergency Obstetric Care in Dhaka's Slums. MANOSHI working paper series no.10.2010; Manoshi WP 10:1-36 published by ICDDR,B, BRAC
18. Gupta PR, Chaudhary SN, Gonnade NV. Maternal and fetal outcome in referred patients to tertiary care centre *Sch J App Med Sci*. 2016; 4(5C):1624-63.
19. Devineni K, Sodumu N. A study of spectrum of referral pattern at a tertiary teaching hospital

- 
- towards better obstetric care. IAIM, 2016;3 (8):193-198.
20. Patel HC, Singh BB, Moitra M, Kantharia SL. Obstetric Referrals: scenario at a primary health centre in Gujarat. Natl J Community Med. 2012;3 (4):711-4.
21. Sabale U, Patankar AM. A Study of Maternal and Perinatal Outcome in Referred Obstetrics Cases. JEvol Med Dent Sci.2015;4(26): 4448-4455.DOI: 10.14260/jemds / 2015/643
22. Charu R, Kamal G, Neelu S. Review of referred obstetric cases–Maternal and Perinatal Outcome. Bombay Hospital J. 2010; 52(1):53.
23. Khatoon A, Hasny SF, Irshad S, Ansari J , An audit of obstetric referrals to Abbasi Shaheed Hospital. Pak J Surg. 2011;27(4):304-8.
24. Poornima M, Rekha G Daver. Maternal and Fetal Outcome in Obstetric Referred Cases. Glob J Res Analys. 2018;7(2):544-7.

**Conflict of Interest: Nil**

**Source of support:Nil**