

Assessment of the depression, anxiety and stress levels among the medical undergraduate students using DASSSanjiw Kumar^{1*}, Amar Kumar²*1,2 Tutor, Department of Community Medicine, Anugrah Narayan Magadh Medical College and Hospital, Gaya, Bihar, India***Received: 15-10-2020 / Revised: 19-11-2020 / Accepted: 10-12-2020****Abstract**

Aims:To assess the depression, anxiety and stress levels among the medical students by using DASS. **Materials and Methods:**This study was carried out in the Department of Community Medicine, Anugrah Narayan Magadh Medical College and Hospital, Gaya, Bihar, India from 1 year, after taking the approval of the protocol review committee and institutional ethics committee. After taking informed consent detailed history was taken from the Participant. They were informed about the anonymous and voluntary nature of participation in the study without any undue fear, stigma, or adverse documentation and were contacted during their free time. A previously validated and standardized survey instrument, Depression Anxiety Stress Scale (DASS 21), was used to collect information on depression, anxiety, and stress. **Results:**A total of 200 students participated in the study giving a response rate of 96%. The profile of the study sample was predominantly male (65%); hosteller (70%) with 61% of students having one sibling. Nearly 25% and 32.5% of students reported having ever smoked or consumed alcohol, respectively. The overall mean age of students was 21.33 (standard deviation=1.98) years. It was also found that 35% had a family history of chronic noncommunicable disease; 12% further mentioned that there was a family history of chronic mental illness while 22% of students had suffered with some medical conditions such as typhoid, malaria, pneumonia, and hospitalization due to injury in the past. It was found that 54(27%), 60(30%), and 86(43%) medical students were affected by symptoms suggestive of depression, anxiety, and stress, respectively. Some students were affected by >1 emotional state. It was observed on bivariate analysis that higher proportion of students with anxiety had a history of some medical condition and this was found to be statistically significant ($P<0.05$). Similarly, family history of chronic noncommunicable disorder was significantly ($P<0.05$) associated with stress and family history of mental illness with depression only. Higher proportion of all the three emotional distress states was found in the 1st-year students in comparison to senior students ($P<0.05$). **Conclusion:** More than half of the medical undergraduate students were found to be affected by depression, anxiety and stress. There is a need for the counselling services to be made available to the students in the medical college to control this morbidity.

Keywords: Depression, Anxiety, Stress, Medical students.

This is an Open Access article that uses a fund-ing model which does not charge readers or their institutions for access and distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>) and the Budapest Open Access Initiative (<http://www.budapestopenaccessinitiative.org/read>), which permit unrestricted use, distribution, and reproduction in any medium, provided original work is properly credited.

Correspondence*Dr. Sanjiw Kumar,**

Tutor, Department of Community Medicine, Anugrah Narayan Magadh Medical College and Hospital, Gaya, Bihar, India.

E-mail: sanjiw00007@gmail.com

Introduction

Worldwide, medical colleges are responsible for making sure that medical students have adequate knowledge and skill before taking professional responsibilities.[1] In order to achieve these goals, medical colleges typically use a curriculum of lectures, simulations supervised practice, mentoring, and hands-on experience to boost individual skill-set. Unfortunately, some aspects of the training process have unintended negative consequences on students' physical and emotional health.[1,2]

Studies revealed that medical students experience a relatively high level of personal distress, with adverse consequences on academic performance, competency, professionalism, and health.[2-5] It is imperative that medical college educator understand the incidences and causes of student distress, adverse consequences on personal and professional well-being, and institutional factors that has an impact on students health.[6] Stress experienced by medical students start from the beginning of the training process.[3] Although some degree of stress is accepted as a normal part of medical training and can be a motivator for some individuals, not all students find the stress manageable.[3-5] Stress may give rise to feelings of fear, incompetence, uselessness, anger, and guilt and has been associated with both psychological and physical disorders.[6] Medical students have used various coping mechanisms to deal with stress; the coping strategies applied by students may determine the effect of stress on psychological and physical health and may determine whether stress has a positive or negative influence. Ineffective stress coping mechanism such as problem avoidance, cynical thinking, social withdrawal, and self-criticism, has negative consequences and can lead to depression, anxiety, and poor mental health. However, appropriate strategies such as problem solving, positive interpretation, and social support can enable students to respond in a manner that leads to adaptation.[6]

In India, the prevalence of depression was around 39%. It was observed that the prevalence of provisionally diagnosed depressive disorder and major depressive disorder in medical students was 21.5% and 7.6%, respectively.[7] Academic performance had a significant association with depression in medical students. The stigma associated with poor academic performance may be a contributing factor. On the other hand, students with excellent academic performance maybe facing pressures due to the competitive nature of medical education.[8] It is no surprise that mental health of medical students in India as an area of

research domain has attracted the second highest attention of the faculty in medical colleges of country after medical education, learning process[9], and evaluation.⁹ With this background, a study was undertaken to assess the prevalence of depression, anxiety, and stress among medical students enrolled in a Patna Medical College, Patna, Bihar, India.

Materials and methods

This study was carried out in the Department of Community Medicine, Anugrah Narayan Magadh Medical College and Hospital, Gaya, Bihar, India from 1 year .after taking the approval of the protocol review committee and institutional ethics committee. After taking informed consent detailed history was taken from the Participant.They were informed about the anonymous and voluntary nature of participation in the study without any undue fear, stigma, or adverse documentation and were contacted during their free time.

Methodology

A previously validated and standardized survey instrument, Depression Anxiety Stress Scale (DASS 21), was used to collect information on depression, anxiety, and stress.[10,11] Additional information was also collected on sociodemographic, academic profile, and personal characteristics of students. Subjective (self) assessment of ability to cope with syllabus and academic performance on a scale of 1–10 points, satisfaction with body image, admission in current medical college, and global satisfaction with life was also assessed. Self assessment scale from 1 to 10 points was classified into low(1–4), medium(5–7), and high(8–10) score. DASS (21 item) is a short scale that allows simultaneous assessment of the three emotional states of depression, anxiety, and stress and each domain contains 7 items, divided into subscales with similar content. The depression scale assesses dysphoria, hopelessness, devaluation of life, self-deprecation, and lack of interest/involvement, anhedonia, and inertia. The anxiety scale assesses autonomic arousal, skeletal muscle effects, situational anxiety, and subjective experience of anxious affect. The stress scale is sensitive to levels of chronic nonspecific arousal. It assesses difficulty relaxing, nervous arousal, and being easily upset/agitated, irritable/over reactive, and impatient. Respondents were asked to document on a 4-point severity/frequency scales to rate the extent to which they have experienced each state over the past week. The scale is as follows: did not apply to me at all = 0;

applied to me to some degree or some of the time = 1; applied to me to a considerable degree or a good part of the time = 2; and applied to me very much or most of the time = 3. Scores for depression, anxiety and stress are calculated by summing the scores for the relevant items and analyzed as per guidelines. The tool is easy to apply in both clinical and nonclinical settings and suitable for use in different age groups including medical students.

Statistical analysis

The data were analyzed using statistical software, SPSS (ver. 20.0) (IBM Inc, Armonk, New York, USA). Descriptive statistics and bivariate and regression analysis were carried out to find association and correlation and considered significant at $P < 0.05$. The internal consistency, i.e., Cronbach's alpha value was 0.87 that was suggestive of high reliability.

Results

A total of 200 students participated in the study giving a response rate of 96%. The profile of the study sample was predominantly male (65%); hosteller (70%) with 61% of students having one sibling. Nearly 25% and 32.5% of students reported having ever smoked or consumed alcohol, respectively. The overall mean age of students was 21.33 (standard deviation=1.98) years. Table 1 depicts sociodemographic profile of study participants. It was also found that 35% had a family history of chronic noncommunicable disease; 12% further mentioned that there was a family history of chronic mental illness while 22% of students had suffered with some medical conditions such as typhoid, malaria, pneumonia, and hospitalization due to injury in the past. It was found that 54(27%), 60(30%), and 86(43%) medical students were affected by symptoms suggestive of depression, anxiety, and stress, respectively. Some students were affected by >1 emotional state. It was observed on bivariate analysis that higher proportion of students with anxiety had a

history of some medical condition and this was found to be statistically significant ($P < 0.05$). Similarly, family history of chronic noncommunicable disorder was significantly ($P < 0.05$) associated with stress and family history of mental illness with depression only. Table 2 shows association of academic variables of students with depression, anxiety, and stress, and it was found that enrollment batch and ability to cope with syllabus were statistically associated with depression, anxiety, and stress ($P < 0.05$). Higher proportion of all the three emotional distress states was found in the 1st-year students in comparison to senior students ($P < 0.05$).

Table 3 depicts additional personal details of medical students affected by emotional state. It was noted that 10.5% reported parental conflict; 17.5% were "always" fearful about future life; 22.5% had poor relationship with family members; 20% were unsatisfied with their body image; and 22% were globally dissatisfied. It was found that satisfaction with body image and global satisfaction with life was statistically ($P < 0.05$) associated with depression and anxiety while relationship with family members was statistically ($P < 0.05$) associated with depression only. Higher proportion of student with depression had fair (poor) relationship with their respective families. Subjective (self) assessment of ability to cope with medical syllabus was inversely but statistically associated ($P < 0.01$) with depression and anxiety, i.e., as the ability to cope with syllabus increases, the probability of occurrence of depression and anxiety decreases. As shown in Table 4, one unit improvement in ability to cope with syllabus results in reduction of 1.36 units in depression and 0.77 units in anxiety. We further studied the correlation between depression, anxiety, and stress and it was found that they were highly correlated with each other. The correlation coefficient value between depression and anxiety was 0.72, depression and stress was 0.73, and anxiety and stress was 0.78.

Table 1: Sociodemographic profile of study participants(n=200)

Variable	N	(%)
Gender		
Male	130	65
Female	70	35
Religion		
Hindu	160	80
Others	40	20
Residence		
Hostel	140	70
Dayscholar	60	30
Number of siblings		

One	122	61
Atleast two	78	39
Students who have ever smoked		
Yes	50	25
Students who have ever consumed alcohol		
Yes	65	32.5

Table 2: Academic profile of study participants affected by the psychological state

Variable	Total (n=200), n (%)	Depression (n=54), n (%)	Anxiety (n=60), n (%)	Stress(n=86), n (%)
Enrollment batch (year of college admission)				
I st year	50 (25)	20 (37.03)*	21 (35)#	31(36.05) ^a
II nd year	55(27.5)	14 (25.92)*	13 (21.67)#	21(24.41) ^a
III rd year	51(25.5)	11 (20.37)*	12 (20)#	18(20.93) ^a
IV th year	44 (22)	9(16.67)*	14 (23.33)#	16 (18.60) ^a
No of attempts to join MBBS				
First attempt	90 (45)	22(40.74)	24 (40)	30 (34.88)
At least 2 attempts	110 (55)	32 (59.26)	36(60)	56 (65.12)
Reason to join MBBS				
Personal choice	170(85)	49 (90.74)	53 (88.33)	74 (86.05)
Parents' pressure	30 (15)	5 (9.26)	7 (11.67)	12 (13.95)
Awareness of vastness of medical course before joining MBBS				
Yes	135 (67.5)	35 (64.81)	44 (73.33)	64 (74.42)
No	65 (32.5)	19 (35.19)	16 (26.67)	22 (25.58)
Number of supplementary examinations				
None	175 (87.5)	47 (87.04)	53 (88.33)#	76 (88.37)
At least one	25 (12.5)	7 (12.96)	7 (11.67)#	10 (11.63)
Satisfaction with regard to admission in this college				
Satisfied	160(80)	40 (74.07)	47 (78.33)	68(79.07)
Unsatisfied	40 (20)	14 (25.93)	13 (21.67)	18(20.93)
Satisfaction with regard to MBBS as a professional carrier				
Satisfied	183 (91.5)	49 (90.74)	56 (93.33)	80(93.02)
Unsatisfied	17 (8.5)	5 (9.26)	4 (6.67)	6 (6.98)
Subjective (self) assessment of ability to cope with medical syllabus on a scale of 1-10 points				
1-4 (low)	20(10)	5 (9.26)*	5 (8.33)#	10 (11.62) ^a
5-7 (medium)	130 (65)	39 (72.22)*	36 (60)#	54 (62.79) ^a
8-10 (high)	50 (25)	10 (18.52)*	19 (31.67)#	22 (25.58) ^a
Subjective (self) assessment of academic performance on a scale of 1-10 points				
1-4 (low)	35 (17.5)	13(24.07)	13 (21.67)	21 (24.42)
5-7 (medium)	134(67)	32 (59.26)	37 (61.67)	53(61.63)
8-10 (high)	31(15.5)	9 (16.67)	10 (16.66)	12(13.95)

*,#, ^a P<0.05

Table 3: Personal profile of study participants affected by emotionalstate

Variable	n (%)	Depression	Anxiety	Stress
	n =200(%)	n =54(%)	N=60(%)	N=86 (%)
History of parental conflict				
Yes	21 (10.5)	6 (11.11)	7 (11.67)	8 (9.30)
Fear of future life				
Always	35 (17.5)	12(22.22)	12 (20)#	11 (12.79)
Sometimes	136 (66)	34 (62.96)	42 (70)#	60(69.77)
Never	29 (14.5)	8 (14.81)	6 (10)#	15 (17.44)
Relationship with friends				
Strong	92(46)	20 (37.04)	26 (43.33)	46(53.49)
Fair	108(54)	34 (62.96)	34 (56.67)	40(46.51)
Relationship with family				
Strong	155 (77.5)	44 (81.48)*	42 (70)	69 (80.23)
Fair	45 (22.5)	10 (18.51)*	18 (30)	17 (19.77)
Satisfaction with body image				
Satisfied	160 (80)	44(81.48)*	47 (78.33)#	61 (70.93)
Not satisfied	40(20)	10(18.51)*	13 (21.67)#	15 (29.07)

^aP<0.05

Table 4: Ordinal regression analysis of association of depression, anxiety, and stress with sociodemographic, academic, and personal profile of medical students

Variable	Estimate	Significance	Significance	95%CI
Depression				
Subjective assessment of ability to cope with medical syllabus	-1.36		0.001	-1.95--0.61
Anxiety				
Subjective assessment of ability to cope with medical syllabus	-0.77		0.01	-1.20--0.18

CI: Confidence interval, LL: Lower limit, UL: Upper limit

Discussion

This descriptive study using DASS-21 scale was suggestive of high prevalence of depression (27%), anxiety (30%), and stress (43%) among medical students enrolled in Anugrah Narayan Magadh Medical college and Hospital, Gaya, Bihar, India. The predictors such as low ability to cope with syllabus and newly entrant students were significantly ($P<0.05$) associated with the emotional distress. This study also reported protective factors such as strong relationship with family members, negative history of medical

condition, satisfaction with body image, and global satisfaction with life. However, a recent large sample survey in southern part of India reported an overall prevalence of depression of 15.9% among general population.[12] In a similar study from Brazil using DASS scale, 34.6%, 37.2%, and 47.1% of medical students suffered from depression, anxiety, and stress, respectively.[13] A study from Turkey found that 27.1% of students were depressed, 47.1% from anxiety, and 27% students were stressed. In a study from Nepal, depression was reported to be 29.9%, anxiety 41.1%,

and stress 27% among medical students.[14] A study based in the United States found 24% of medical students to be depressed while another study from the US reported 12% of medical students to be diagnosed with probable major depression using DSM III criteria.[15] In a study from Egypt, 43.9% of students were suffering from anxiety.[16] Similar alarming statistics have been corroborated by our study also. Studies conducted in various regions of India reflect diverse situation depending on the use of study instruments. In a study from Bhubaneswar (Odisha), the prevalence of depression, anxiety, and stress among medical students was 51.3%, 66.9%, and 53%, respectively, using DASS scale.[17] Another study reported that 39.44% of students suffered from depression, 66.05% from anxiety, and 51.37% from stress.[18] A Jodhpur (Rajasthan)-based study found that 57.98% of students depressed and 47.41% suffered from anxiety.[19] Another Delhi-based study found the overall prevalence of provisionally diagnosed depressive and major depressive disorders among medical students to be 21.5% and 7.6%, respectively.[20] On the contrary, some studies conducted two decades ago have found little or no evidence of stress among medical students.[21,22] On joining medical college, students embark their professional journey with high expectations and are loaded with lots of new information to be crammed which at times become difficult to process. The students leave the protected, pampered, and very supportive environment of their family and come to stay in hostel under highly competitive environment. This could be contributing to the higher prevalence of depression, anxiety, and stress seen in 1st-year medical students in this study. Similar observation was reported by Aktekin regarding worsening in overall mental health with high level of depression and anxiety among 1st- and 2nd- year medical students.[23] Quince et al. performed a longitudinal study at a UK medical school and found a prevalence of depression ranging from 5.7% to 10.6% among students in the basic years and 2.7% to 8.2% in students from clinical stages of the course.[24] Further, there was no significant difference of emotional state between males and females in our study. In our urban-based study, 20% of students were unsatisfied with their body image which is higher than another study conducted at Rohtak, Haryana (13.5%), with a predominantly rural flavor.[25] The content of MBBS subject in medical colleges of India is based on global best practices, and course duration is of four-and-half years followed by 1-year internship. The students are systematically and in structured way exposed to different subject streams (preclinical,

paraclinical, and clinical) through processes prescribed under regulatory body with student to keep record of daily activity/learning in a log book followed by formative and summative evaluation.[26] It is no surprise that India is one of the popular destinations of medical, transplant, reproductive, and health tourism in the world, and a substantial proportion of renowned doctors working in developed country have roots in this great nation.[27,28] It could be partly attributed to rigor of professional training and the quality of learning environment in medical colleges of India. The development and growth potential of an individual is directly correlated with the depth of knowledge, positive attitude, and diverse skills she/he possesses and demonstrates whether at undergraduate or higher level.[29] It can be though attributed to early start and purposeful planning but also indicating a case scenario of intense peer pressure, uncertain future environment, rising stress, and anxiety. Medical students under strain are either unaware of their situation or reluctant to seek help. High-risk students found in our study were provided personal and confidential counseling under additional supervision of mental health expert.

Conclusion

More than half of the medical undergraduate students were found to be affected by depression, anxiety and stress. There is a need for the counselling services to be made available to the students in the medical college to control this morbidity.

References

1. Liaison Committee on Medical Education (LCME). Functions and structure of a medical school: standards for accreditation of medical education programs leading to the M.D. degree. 2003.
2. Association of American Medical Colleges (AAMC). Report I. Learning objectives for medical student education: guidelines for medical schools. Medical schools objectives project. 1998.
3. Abdulghani HM, AlKanhil AA, Mahmoud ES, Ponnampereuma GG, Alfaris EA. Stress and its effects on medical students: a cross-sectional study at a college of medicine in Saudi Arabia. *J Health Popul Nutr.* 2011; 29(5):516.
4. Abdulghani HM, Irshad M, Al Zunitan MA, Al Sulihem AA, Al Dehaim MA, Al Esefir WA et al. Prevalence of stress in junior doctors during their internship training: a cross-sectional study of three Saudi medical colleges' hospitals. *Neuropsychiatr Dis Treat* 2014;10:1879–1886.
5. AlFaris EA, Naeem N, Irfan F, Qureshi R, van der Vleuten C. Student centered curricular elements are associated with a healthier educational environment

- and lower depressive symptoms in medical students. *BMC Med Educ*.2014; 14(1):192.
6. Dyrbye LM, Thomas MR, Shanafelt TD. Medical student stress: causes, consequences, and proposed solutions. *Mayo Clin Proc*.2005; 80(12):1613–1622.
 7. Iqbal S, Gupta S, Venkatarao E. Stress, anxiety & depression among medical undergraduate students & their socio-demographic correlates. *Indian J Med Res* 2015;141(3):354-8.
 8. Sidana S, Kishore J, Ghosh V, Gulati D, Jiloha RC, Anand T. Prevalence of depression in students of a medical college in New Delhi: a cross-sectional study. *Australasian Med J* 2012;5(5):247-50.
 9. Sachdeva S, Sachdev TR, Sachdeva R, Dwivedi N, Taneja N. Published research studies conducted amongst Indian medical undergraduate students: bibliometric analysis. *Indian J Community Health* 2017;29.
 10. Lovibond SH, Lovibond PF. *Manual for the Depression Anxiety and Stress Scales*. 2nd ed. Sydney: Psychology Foundation; 1995.
 11. Crawford JR, Henry JD. The depression anxiety stress scales (DASS): Normative data and latent structure in a large non-clinical sample. *Br J ClinPsychol* 2003;42:111-31.
 12. Poongothai S, Pradeepa R, Ganesan A, Mohan V. Prevalence of depression in a large urban South Indian population – The Chennai urban rural epidemiology study (CURES-70). *PLoS One* 2009;4:e7185
 13. IL Damasio MO, Natalia De C, Maddalena P, Kleinsorge R, GraneroLucchetti AL, Helena Cerrato S, et al. Depression, stress and anxiety in medical students: A cross-sectional comparison between students from different semesters. *Rev Assoc Med Bras* 2017;63:1-5.
 14. Kunwar D, Risal A, Koirala S. Study of depression, anxiety and stress among the medical students in two medical colleges of Nepal. *Kathmandu Univ Med J (KUMJ)* 2016;14:22-6.
 15. Givens JL, Tjia J. Depressed medical students' use of mental health services and barriers to use. *Acad Med* 2002; 77:918-21.
 16. Zoccolillo M, Murphy GE, Wetzel RD. Depression among medical students. *J Affect Disord* 1986 ;11:91-6.
 17. Iqbal S, Gupta S, Venkatarao E. Stress, anxiety and depression among medical undergraduate students and their socio-demographic correlates. *Indian J Med Res* 2015;141:354-7.
 18. Vaidya PM, Mulgaonkar KP. Prevalence of depression anxiety and stress in undergraduate medical students and its correlation with academic performance. *Indian J OccupTher* 2007;39:7-10.
 19. Hakim A, Tak H, Nagar S, Bhansal S. Assessment of prevalence of depression and anxiety and factors associated with them in undergraduate medical students of Dr. S. N. Medical College, Jodhpur. *Int J Community Med Public Health* 2017;4:3267-7
 20. Sidana S, Kishore J, Ghosh V, Gulati D, Jiloha R, Anand T, et al. Prevalence of depression in students of a medical college in New Delhi: A cross-sectional study. *Australas Med J* 2012;5:247-50.
 21. Bramness JG, Fixdal TC, Vaglum P. Effect of medical school stress on the mental health of medical students in early and late clinical curriculum. *ActaPsychiatrScand* 1991;84:340-5.
 22. Vaz RF, Mbajorgu EF, Acuda SW. A preliminary study of stress levels among first year medical students at the University of Zimbabwe. *Cent Afr J Med* 1998;44:214-9.
 23. Aktekin M, Karaman T, Senol YY, Erdem S, Erengin H, Akaydin M, et al. Anxiety, depression and stressful life events among medical students: A prospective study in Antalya, Turkey. *Med Educ* 2001;35:12-7.
 24. Quince TA, Wood DF, Parker RA, Benson J. Prevalence and persistence of depression among undergraduate medical students: A longitudinal study at one UK medical school. *BMJ Open* 2012;2. pii: e001519.
 25. Goswami S, Sachdeva S, Sachdeva R. Body image satisfaction among female college students. *Ind Psychiatry J* 2012;21:168-72.
 26. Sachdeva R, Sachdeva S. Medical education, training and patient care from the lens of resident. *Natl J Community Med* 2012;3:750-3.
 27. Jose R, Sachdeva S. Keeping an eye on future: Medical tourism. *Indian J Community Med* 2010;35:376-8.
 28. Sulania A, Sachdeva S, Jha D, Kaur G, Sachdeva R. Organ donation and transplantation: An updated overview. *MAMC J Med Sci* 2016;2:18-27.
 29. Sachdeva S, Sachdev TR. Skills and practices for the postgraduate trainees of community medicine, public health, and hospital administration courses in India: Learn to demonstrate and imbibe. *J SciSoc* 2016;43:109-11.

Conflict of Interest: Nil

Source of support:Nil