

An observational study of assessment of pulmonary function and clinical features, and their comparison between obese and non-obese patients of bronchial asthma

K.Rajasekhar*

Assistant Professor, Department of Pulmonary Medicine, Maheswara Medical College & Hospital, Hyderabad, Telangana, India

Received: 23-10-2020 / Revised: 27-11-2020 / Accepted: 13-12-2020

Abstract

Introduction: Asthma is a heterogeneous, inflammatory disorder of airways and is associated with airway hyper responsiveness that leads to recurrent episodes of wheezing, breathlessness, chest tightness and coughing particularly at night or in the early morning. These episodes are usually associated with wide spread but variable airflow obstruction within the lung that is often reversible either spontaneously or with treatment. The incidence of obesity has doubled worldwide since 1990. In India 12.1% males and 16% females are obese. **Materials and Methods:** Patients treated under Department of Pulmonary Medicine, attached to Maheswara Medical College & Hospital, Telangana, Hyderabad, with bronchial asthma, fulfilling the inclusion & exclusion criteria were included in the study after obtaining written informed consent. Demographic data, history, clinical examination and details of investigations like Pulmonary function test, Complete hemogram (Hemoglobin, Total leukocyte count and Differential leukocyte count), Chest X-ray (Postero-anterior view), Electrocardiogram, Lipid profile -Triglyceride, Total cholesterol, High density lipoprotein (HDL) and low density lipoprotein (LDL), Sputum AFB, Renal function test. Sample size taken for convenience. **Results:** In this study, out of 100 patients 55% were males, and 45% were females with higher total cholesterol and triglycerides in obese asthmatics compared to non-obese asthmatics. The most common symptoms were breathlessness followed by cough and wheeze. In this study, it was found that breathlessness, cough, wheeze are more common in obese asthmatics than non-obese asthmatics. There was significant difference of waist circumference (W.C.), hip circumference (H.C.) and waist to hip ratio (W.H.R.) between obese and non-obese asthmatics. Hypertension and diabetes mellitus more prevalent in obese group than non-obese. FVC, FEV1, and FEF25-75 % values were reduced in obese asthmatics compared to non-obese asthmatics in contrast to FEV1/FVC which was higher in obese asthmatics compared to non-obese asthmatics. It was observed that increased in BMI causes impaired pulmonary function. **Conclusion:** The increasing prevalence of asthma and obesity has suggested an association between the two. The most common symptoms observed in this study were breathlessness, cough and wheeze. PR (Pulse Rate) SBP (Systolic Blood Pressure), DBP (Diastolic Blood Pressure), RR (Respiratory Rate), TLC (Total Leucocyte Count), AEC (Absolute Eosinophil Count), LP (Lipid Profile) were higher in obese asthmatics. Diabetes mellitus and hypertension were more prevalent in obese asthmatics. FVC, FEV1 are decreased in obese asthmatics but the amount of reversibility is more for non-obese asthmatics. It was also observed that increased BMI causes impaired pulmonary function.

Keywords: Asthma, HDL, LDL, FVC, FEV1

This is an Open Access article that uses a fund-ing model which does not charge readers or their institutions for access and distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>) and the Budapest Open Access Initiative (<http://www.budapestopenaccessinitiative.org/read>), which permit unrestricted use, distribution, and reproduction in any medium, provided original work is properly credited.

Introduction

Asthma is a heterogeneous, inflammatory disorder of

airways and is associated with airway hyper responsiveness that leads to recurrent episodes of wheezing, breathlessness, chest tightness and coughing particularly at night or in the early morning. These episodes are usually associated with wide spread but variable airflow obstruction within the lung that is often reversible either spontaneously or with treatment[1]. The incidence of obesity has doubled

*Correspondence

Dr. K Rajasekhar,

Assistant Professor, Department of Pulmonary Medicine, Maheswara Medical College & Hospital, Hyderabad, Telangana, India.

E –Mail: haasinirajajuly3@gmail.com

worldwide since 1990[2]. In India 12.1% males and 16% females are obese[3]. Asthma in the obese is more difficult to control[4] and is associated with poor quality of life and requires more health care resources than in non-obese asthmatics[2]. Respiratory symptomatology is more in obese asthmatics because of alterations in chest wall mechanics and also because of associated comorbidities like hormonal imbalance, GERD, increased truncal fat and decreased exercise tolerance. Despite these observations, obesity alone has not been shown to cause dyspnea in persons at rest[3]. Bronchial asthma is diagnosed based on clinical features and spirometry. There is decreased FEV₁, decreased FEV₁/FVC and reversibility is more than 12% and 200ml after 15 minutes of inhaled bronchodilator administration. For the convenience of management GINA (Global Initiative for Asthma) has classified asthma into the following categories[4-6].

- Intermittent (Symptoms <once a week, FEV₁ or PEF ≥ 80% of predicted).
- Mild Persistent (Symptoms >once a week <once a day FEV₁ or PEF ≥ 80% of Predicted).
- Moderate Persistent (Symptoms daily FEV₁ or PEF 60 to 80% of predicted).
- Severe Persistent (Symptoms daily FEV₁ or PEF <60 % predicted).

Materials and methods

Patients treated under Department of Pulmonary Medicine, attached to Maheswara Medical College & Hospital, Telangana, Hyderabad, with bronchial asthma, fulfilling the inclusion & exclusion criteria were included in the study after obtaining written informed consent. Demographic data, history, clinical examination and details of investigations like Pulmonary function test, Complete hemogram (Hemoglobin, Total leukocyte count and Differential leukocyte count), Chest X-ray (Postero-anterior view), Electrocardiogram, Lipid profile -Triglyceride, Total cholesterol, High density lipoprotein (HDL) and low density lipoprotein (LDL), Sputum AFB, Renal function test. Sample size taken for convenience.

Results

Table 1: WHO Classification of BMI (2003)

Classification	BMI (Kg/m ²)	Risk of Comorbidities
Underweight	Less than 18.5	Low-but risk of other clinical problems increased
Normal Range	18.5-24.9	Average
Over weight	>25	
Pre Obese	25-29.9	Increased
Obese Class I	30-34.9	Moderate
Obese Class II	35-39.9	Severe
Obese Class III	>40	Very Severe

Source of Data

The study was conducted in Department of Pulmonary Medicine, Maheswara Medical College & Hospital, Telangana, Hyderabad on patients with Stable obese and non-obese Bronchial Asthma patients who were diagnosed with bronchial asthma as per GLOBAL Initiative for Asthma (GINA) guidelines undergoing pulmonary function test on inpatient or outpatient basis.

Study Design: A Hospital based cross sectional observational study.

Place of Study: Maheswara Medical College & Hospital, Telangana, Hyderabad.

Duration of Study: One year (January 2019 – December 2019).

Sampling Technique: Convenient sampling.

Sample Size: A total of 100 study subjects-50 obese and 50 non-obese asthmatic adults. BMI <22.9 is taken as Non-Obese and >23 as Obese as per WHO Asia Pacific perspective for Asians (WHO IOTF 2003)10 Inclusion Criteria1. All individuals above 18 years of age.2.All adults diagnosed with Bronchial asthma as per GINA1guidelines.3.All asthmatic adults who are obese (which included at risk, Obese I and Obese II) and non-obese (underweight and normal) as per WHO Asian pacific perspective for Asians.4.Patients who are willing to participate. Exclusion Criteria1Unstable inpatients/out-patients. (i.e. Acute exacerbation of asthma)2Patients with chronic obstructive pulmonary disease, Bronchiectasis and other respiratory diseases.3Patients with malignancies, cardiac complications or neuromuscular diseases.4Post-operative patients, individuals with serious systemic illnesses like chronic renal failure or complicated diabetes mellitus etc.

Statistical Methods: Data was entered in Microsoft excel and analyzed using EPI INFO version 6 software. Descriptive statistics like proportions, percentages and standard deviation are used. For comparison between the groups chi square test, Independent 't' test, ANOVA were used. P value of less than 0.05 is considered as statistically significant.

Table 2: WHO Asia Pacific Perspective for Asians (WHO IOTF 2003)

Classification	BMI (Kg/m ²)	Risk of Comorbidities
Underweight	Less than 18.5	Low-but risk of other clinical problems increased
Normal Range	18.5-24.9	Average
Over weight	>25	
At Risk	25-29.9	Increased
Obese Class I	30-34.9	Moderate
Obese Class II	35-39.9	Severe

Table 3: Comparing PRE FVC, FEV1, FEV1/FVC, FEF (25-75) in Obese and Non-Obese Asthmatics

PFT Values	Obese			Non Obese		
	Predicted	Observed	% of Observed FVC	Predicted	Observed	%
FVC	2.44 +0.49l/s	1.61 +0.38 l/s	66.87±13.69%	2.82 +0.73l/s.	2.26+0.88l/s.	79.76±22.04%
FEV1	2.10 +0.40.	1.13 +0.24	55.04±11.35%	2.47 +0.66	1.50 +0.65.	59.72±18.80%
FEV1/FVC	86.10 +4.84	70.40 +8.01	86.10 +4.84%	87.08 +4.42	65.03 +7.34.	87.08+4.42%
FEF(25-75)	3.20 +0.55	1.31 +0.62	42.23±21.05%	3.87 +0.78	1.59 +0.76	40.80±17.14%

Table 4: Comparing POST FVC, FEV1, FEV1/FVC, FEF (25-75) in Obese and Non-Obese Asthmatics

PFT Values	Obese		Non Obese	
	Observed	% of observed	Observed	% of observed
FVC	1.92 +0.45 l/s	79.58 ± 13.83 %	2.56+0.80 l/s.	91.65 ± 21.26 %
FEV1	1.47 +0.33	70.85 ± 13.04 %	1.94 +0.72	78.20 ± 21.26 %
FEV1/FVC	77.07 +7.92	-	74.38 +9.41	-
FEF (25-75)	1.93+0.68	61.35 ± 21.01 %	2.76 +1.01	71.92 ± 24.00 %

A total of 100 bronchial asthma patients diagnosed as per GINA guidelines were taken up for the study. Out of 100 patients, 50 were obese bronchial asthma patients (n =25) and remaining 50 were non-obese bronchial asthma patients (n=25). In this study of 100 patients, the age of patients ranged from 20-59 years, the age of patients in obese group ranged from 28-59 years with a mean of 40.27+8.395 years, in non-obese group it ranged from 20-50 years with a mean of 32.57+7.583 years. Out of total 100 asthmatic patients studied, in obese asthmatic group 18(36.6%) were male patients and 32 (63.4%) were female patients and in non-obese asthmatic group 36 (73.3%) were males and 14 (26.7%) were females. The association between the groups is statistically significant. (p=0.009)

Hypertension-Out of 100 patients studied, 15 (30%) obese asthmatics and 4(6.66%) non obese asthmatics had previous history of hypertension. There was significant association seen between the groups (p=0.02)b Diabetes Mellitus-Out of 100 patients studied, 23.3% obese asthmatics and 6.66% non-obese asthmatics had previous history of diabetes mellitus. The mean percentage change in reversibility of spirometric values in non-obese population is 34.07%

whereas in obese population it is 29.24%.The student t test applied found that the p value between the groups for this variable is not statistically significant. (p=0.23776) Bronchodilator response is found to be more in non-obese asthmatic group compared to obese asthmatics, however it was not statistically significant.

Discussion

In this study of 100 patients with bronchial asthma, the age of patients ranged from 20-59 years, the mean age of patients in obese group was 40.27 +8.395years, in non-obese group was 32.57+7.583 years. Nearly two thirds in obese group were female and male in non-obese group. The mean weight of the patients in the obese group was 72.53 +12.822 kgs. and in the non-obese group with a mean of 54.13 +9.464 kgs. Castro-Rodriguez et al.,14(2001) demonstrated that girls becoming overweight or obese between 6 and 11 years of age had increased odds of developing new asthma symptoms. The mean BMI of the patients in the obese group was 30.16 +4.63 kilograms/metre² and in the non-obese group was 20.55 +2.689 kg/m². The study findings is similar to the study done by Dosi R et

al[9].Beuther et al (2007)[10] have demonstrated a clear dose response relationship between BMI and asthma, suggesting that asthma risk increases further as body weight increases. In addition, he had shown that the odds of incidence of asthma in overweight and obese men and women were similar. Most common symptom in both groups was breathlessness which was present in all patients in obese group and 90% of non-obese group, which was followed by cough and wheeze. The present study showed significant association of cough and seasonal variation between obese and non-obese asthmatic patients as comparable with the study done by Aruna G et al[11] The presence of comorbid condition like diabetes and hypertension was more common in obese compared to non-obese which was consistent with the study by Pakhale S et al (2010)[12].Our study shows there is significant difference of Absolute eosinophil count between the groups. ($p=0.05$) but it was contrast with the finding of the study done by Aruna G et al[11].The mean of Forced expiratory volume in 1st second (FEV1) and Forced vital capacity (FVC), in obese asthmatics is 1.13 ± 0.24 l/s and 1.61 ± 0.38 l/s respectively and in non-obese it is 1.50 ± 0.65 l/s and 2.26 ± 0.88 l/s. In the present study FEV1, FVC is lower in obese asthmatics compared to non-obese asthmatics which is comparable with the studies done by Pakhale s et al.(2010) [12]Dosi R et al, Razi et al[13].The mean percentage change of reversibility in spirometric values (FEV1) among non-obese population is 34.07%, whereas in obese population it is 29.24%.but was not statistically significant. ($p=0.23776$) which was in contrast with the study done by Sharma L, et al [14]which showed highly significant difference found in the spirometric variables in obese and non-obese asthmatics. Our study also compared mean value of FEV1/FVC % observed for the obese and non-obese groups which was 70.40 ± 8.01 and 65.03 ± 7.34 respectively. This finding is similar to the study done by Sharma et al[14] whereas it is contrast with the findings of the study done by Dosi et al[9]. The mean of FEF 25-75% in the present study among obese asthmatics is 1.31 ± 0.62 l/s and in non-obese it is 1.59 ± 0.76 l/s. The values were lower compared to non-obese asthmatics which was comparable with the study done by Zied et al.Earlier studies showed (Sharma L, et al)[14] the mean percentage change in reversibility of spirometric values in non-obese population highly significant, but our study shows the contrast[9].

Conclusion

The increasing prevalence of asthma and obesity has suggested an association between the two. The most

common symptoms observed in this study were breathlessness, cough and wheeze. PR (Pulse Rate) SBP (Systolic Blood Pressure), DBP (Diastolic Blood Pressure), RR (Respiratory Rate), TLC (Total Leucocyte Count), AEC (Absolute Eosinophil Count), LP (Lipid Profile) were higher in obese asthmatics. Diabetes mellitus and hypertension were more prevalent in obese asthmatics. FVC, FEV1 are decreased in obese asthmatics but the amount of reversibility is more for non-obese asthmatics. It was also observed that increased BMI causes impaired pulmonary function.

References

1. From the Global Strategy for Asthma Management and Prevention. Global Initiative for Asthma 2014 available from <http://www.ginaasthma.org/>.
2. Centres for Disease Control and Prevention obesity trends in the U.S <http://www.cdc.gov/ncd/php/obesity/trend/index.htm>.
3. National Family Health survey, 2005–06 Mumbai International Institute for Population Sciences, 2007.
4. Mosen DM, Schatz M, Magid DJ et al. The relationship between obesity and asthma severity and control in adults J. Allergy Clin. Immunol. 2008; 122: 507-11.
5. Stanley S., Rogers M. Obesity as a cause of dyspnea in otherwise healthy men. Am. Family Physician, 1999 ;ch 59 : 1280-86
6. Global initiative for Asthma (GINA), National Heart, Lung and Blood Institute (NHLBI), global strategy for asthma management and prevention. Bethesda MD, 2006 p.339. Also at www.ginaasthma.org2014
7. The international classification of adult underweight, over weight and obesity. WHO1995, 2000, 2004
8. Misra A, Khurana L. Obesity and metabolic syndrome in developing countries. J. Clin Endocrinol Metab 2008; 93 (11 suppl.):9
9. Dosi R, VeerapaneniVR, Bhaskar BV, et al. A study of spirometry in obese and non-obese asthmatics. Journal of Evolution of Medical and Dental Sciences 2013; 2(8): 903- 20.
10. Beuther DA, Sutherland ER. Overweight, obesity and incident asthma: a meta-analysis o prospective epidemiologic studies.Am J Respir Crit Care Med 2007;175(7):661-6.
11. Aruna G, Kumar KS, Chandra N. Comparison of prevalence, clinical presentation and spirometry in bronchial asthma in obese

-
- and non-obese patients. Journal of Evolution of Medical and Dental Sciences 2015;4(58):10136-41.
12. Pakhale S, Doucette S, Vandemheen K, et al. A comparison of obese and nonobese people with asthma. Chest 2010;137(6):1316-23.
13. RaziE, MoosaviGA. The effect of positions on spirometric values in obese asthmatic patients. Iran J Allergy Asthma & Immunol 2007;6(3):151-4.
14. Sharma L, Putti N, Chaskar A, et al. To evaluate the influence of inhaled bronchodilator therapy on spirometric test variable in obese and non-obese asthmatics. Journal of medical and scientific research 2015; 3(4): 166-71.

Conflict of Interest: Nil

Source of support: Nil