

Evaluation of breast lesion by ultrasonography during pregnancy and lactating women**Deepak K S¹, Bharat M.P^{1*}, Nandan Kumar LD²**¹ Associate Professor, Department of Radiodiagnosis, Subbaiah Institute of Medical Sciences, Shimoga, Karnataka, India² Assistant Professor, Department of Radiodiagnosis, Subbaiah Institute of Medical Sciences, Shimoga, Karnataka, India

Received: 14-10-2020 / Revised: 12-11-2020 / Accepted: 16-12-2020

Abstract

Introduction: Substantial physiological changes occur during pregnancy and lactation, making breast evaluation challenging in these patients. The aim of this study was to characterize the breast lesion seen in Indian women and to establish the correlation between clinical features of breast lesion and their pathological diagnosis. **Materials and Methods:** This is a prospective and observational study conducted at Subbaiah Institute of Medical Sciences attended between April 2019 to March 2020. All female Patients irrespective of age who came for ultrasound breast examination with suspected breast lesions, complain of pain, palpable lump, breast complaints like nipple discharge, retraction, skin thickening were included in the study. Sixty pregnant or /and lactating females who came for ultrasound examination were done such as proper application of compression, transducer positioning, and image labelling. **Results:** In our present study 100 patients were examined amongst which 39 cases were benign 3 were malignant lesions and rest of the cases were normal. Of the remaining consecutive 39 cases (BIRADS 2 to 6) were evaluated clinically, by sonography and FNAB. In this study, the age range of patients presenting with breast lesions was 18–70 years with mean age 48.3 years. 19 patients were breast lesions occurred in the age group of 40-60 years, 6 were younger than 40 years and 14 were between 61-70 years. Most lesions were found in the upper outer quadrant and the right breast was affected more than the left. 15 lesions were assigned BIRADS 5 category. The second most common category was BI-RADS 2 in 11 lesions. There were total of 39 benign cases, Fibroadenoma (13) being the commonest, followed by Fibrocystic disease (9), Cysts (7), Mastitis (4), Duct ectasia (3), least being Phylloides, Galactocele and Papilloma. **Conclusion:** Evaluation of the pregnant and lactating patients who present with a breast problem is challenging. Although ultrasound may characterise the finding in many cases, mammography and even MRI may have a role in the management of these patients.

Keywords: Breast lesion, Ultrasonography, Pregnant, Lactating women.

This is an Open Access article that uses a fund-ing model which does not charge readers or their institutions for access and distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>) and the Budapest Open Access Initiative (<http://www.budapestopenaccessinitiative.org/read>), which permit unrestricted use, distribution, and reproduction in any medium, provided original work is properly credited.

Introduction

Radiological evaluation varies depending upon the age of the woman, her pregnancy and lactational status. Subsequent to a clinical history and thorough physical examination, patients are frequently imaged to

determine whether there is an underlying abnormality to account for the patient's symptoms. [1] For pregnant and lactating women under the age of 30 years, ultrasound is the initial imaging test of choice given the lack of radiation exposure. Mammogram could be considered in these patients if ultrasound is negative or it reveals indeterminate, suspicious or no findings. [2] Lactating women over 30 years of age are typically imaged using both mammography and ultrasound. In an effort to reduce the overall breast density, lactating patients are encouraged to express milk immediately prior to imaging. In a pregnant patient, mammography

*Correspondence

Dr. Bharat MP

Associate Professor, Department of Radiodiagnosis, Subbaiah Institute of Medical Sciences, Shimoga, Karnataka, India.

E-mail: mpbharat80@gmail.com

should be performed, if ultrasound reveals a suspicious finding or if biopsy of a solid lesion reveals malignancy. A complete evaluation of a pregnant patient with a lump should not be delayed until after delivery, because of fear of radiation. [3]The imaging appearance on ultrasound is variable depending upon the duration of pregnancy and/or lactating state. An overall diffuse increase in breast density accompanied by breast enlargement is commonly seen on mammography. Given increased density of the breast the sensitivity of mammography is low (30 % for dense breast compared with 80 % for fatty breast), and cancer detection may be somewhat difficult. [4] According to one study evaluated patients with false-negative mammograms and symptomatic cancer, and found that 78 % of the mammographically occult lesions were in women with heterogeneously or extremely dense tissue. The imaging features of breast cancer on the mammogram are identical to those seen in non-pregnant women. [5] These are speculated or irregular masses, pleomorphic linear branching or grouped microcalcifications, focal asymmetries and architectural distortion. Detection is sometimes difficult as the overlying dense tissue may obscure the findings. Ultrasound has a better sensitivity in pregnant and lactating patients, ranging from 86.7 to 100 %. [6] On ultrasonography, during pregnancy, the breast shows diffuse hypoechogenicity with fibroglandular enlargement and increased vascularity. In lactating women, the breast shows diffuse hyperechogenicity with a prominent ductal system and increased vascularity. [7] Ultrasonography is the best imaging modality to evaluate breast lesions during pregnancy and lactation, as it is sensitive and confers no radiation exposure. [8]

According to the ACR Appropriateness Criteria, pregnant woman with palpable masses or pathological nipple discharge should be initially evaluated by ultrasonography in order to characterize the features of the lesion and plan proper management. In this article, we discuss and illustrate the pathologic changes that cause most radiologic and cytopathologic diagnostic difficulties during pregnancy and lactation. In addition, we assess medicolegal issues related to pregnancy, particularly the use of mammo-graphy during pregnancy and the risk it poses to the fetus. We emphasize the value of ultrasonography (US) as the most appropriate and effective method of evaluating breast disorders during pregnancy and lactation.

Materials and Methods

This is a prospective and observational study conducted at Subbiah institute of Medical sciences attended between April 2019 to March 2020.

Inclusion criteria

All female Patients irrespective of age who came for ultrasound breast examination with suspected breast lesions, complain of pain, palpable lump, breast complaints like nipple discharge, retraction, skin thickening were included in the study.

Exclusion criteria

Patients with previous history of incision and drainage, already diagnosed or on medical treatment were excluded from the study.

Sixty pregnant or /and lactating females who came for ultrasound examination were done such as proper application of compression, transducer positioning, and image labelling. In order to stabilize, center, and thin out the breast tissue, the conventional position for breast US examination places the patient supine with the arm of the side being examined raised above the head. With larger breasts, a degree of elevation under the shoulder blade may be required in order to center the breast. This can best be accomplished with a foam wedge or roll of towels or sheets. Scanning is performed with the degree of compression necessary to penetrate to the area of interest and eliminate superficial artifact. Scanning in the plane of ductal anatomy can be achieved by scanning in the radial and antiradial planes. Radial scanning is performed with the long axis of the transducer oriented along the long axis of the ductal and lobar anatomy (nipple to periphery of the breast in a branching pattern) and antiradial in the orthogonal plane (from the periphery of the breast inward toward the nipple). Transverse and sagittal plane scanning are acceptable in the initial survey, and if a lesion is detected, radial and antiradial scanning are recommended, as the margins and extension of the mass may be better displayed and this approach increases the potential for finding other masses within the same ductal system. The position of the lesion should be labeled on the image according to the mammographic clock, noting distance from the nipple or areolar margin. It is important that this information be labeled on each image to ensure appropriate follow-up or localization for biopsy. US findings were categorized according to the Breast Imaging Report and Data System (BI-RADS) lexicon using the following tumour classification, [11]: Shape (oval, round or irregular), orientation (parallel to the skin surface or not), margin (circumscribed or not, indistinct, angular, spiculated or microlobulated), echo pattern (anechoic, hypoechoic, hyperechoic or

complex), posterior acoustic features (none, enhancement or shadowing), surrounding tissue change (absent or present), vascularity (none, focal or penetrating flow, or diffusely increased flow), presence of associated calcifications (none or microcalcifications in or out of a mass). The BI-RADS lexicon requires the breast imaging report be summarized into 1 of 7 possible categories, [12]

BI-RADS 0 - Further assessment required;

BI-RADS 1 - Negative study;

BI-RADS 2- Benign finding (risk of malignancy similar to that of the surrounding parenchyma);

BI-RADS 3 - Probably Benign finding (less than 2% risk of malignancies should be followed up at 6, 12, and 24 months, and then classified as benign category 2 after showing stability for 24 months or biopsied if concerning changes or growth are seen);

BI-RADS 4- lesion is Suspicious for Malignancy (biopsy is offered);

BI-RADS 5- lesions are Highly suggestive of Malignancy; and

BI-RADS 6- lesions are Biopsy-proven Malignant before surgery is obtained (it is suggested that appropriate actions should be taken for these categories).

“Positive” category was all those who had BIRADS assessment category 4, 5 and 6. “Negative” category was all those with BIRADS assessment category 0, 1, 2 and 3. All patients underwent pathological assessment either by FNAC or biopsy or both. Biopsies included either core biopsy or surgical excision biopsy. Surgical specimens had been fixed in 10% formaldehyde solution and cut into serial 5-mm thick slices. Histopathological slides in each tumor were reviewed by a pathologist independently. The cytology reports were classified as benign, suspicious for carcinoma, malignant, or inadequate. Histology was performed if cytology was suspicious or suggestive of malignancy. BI-RADS criteria combined with US guided FNAB, were correlated with pathological findings to determine the Sensitivity, Specificity and Accuracy of the Sonographic examinations. After the pathological assessment patients were reviewed in the out-patient’s clinic with their pathology report to plan any further treatment.

Results

In table 1, the present study 100 patients were examined amongst which 39 cases were benign 3 were malignant lesions and rest of the cases were normal.

Table 1: BIRADS Score

BIRADS Score	Age group in years		
	18-40	41-60	61-70
0			
1	0	0	0
2	3	6	2
3	1	3	5
4	1	1	2
5	1	9	5
6	0	0	0
Total	6	19	14

In table 1, of the remaining consecutive 39 cases (BIRADS 2 to 6) were evaluated clinically, by sonography and FNAB. In this study, the age range of patients presenting with breast lesions was 18–70 years with mean age 48.3 years. 19 patients were breast lesions occurred in the age group of 40-60 years, 6 were younger than 40 years and 14 were between 61-70 years. Most lesions were found in the upper outer quadrant and the right breast was affected more than the left. 15 lesions were assigned BIRADS 5 category. The second most common category was BI-RADS 2 in 11 lesions.

Table 2: Sonographic diagnosis of Carcinoma Breast compared with pathologic findings

Sonography	Pathology		Total
	Positive	Negative	
Positive	19	3	22
Negative	2	15	17
Total	21	18	39

The patients with malignant disease underwent surgery. The surgical and histo-pathological findings were positive for carcinoma breast in 21 patients in table 2.

Table 3: Distribution of Benign lumps

Benign lumps	Number of patients
Fibroadenoma	13
Fibrocystic disease	9
Cysts	7
Mastitis	4
Duct ectasia	3
Phylloides	1
Galactocele	1
Papilloma	1

There were total of 39 benign cases, Fibroadenoma (13) being the commonest, followed by Fibrocystic disease (9), Cysts (7), Mastitis (4), Duct ectasia (3), least being Phylloides, Galactocele and Papilloma in table 3.

Table 4: Results of Sonographic studies in diagnosis of Breast Disease

Breast disease	Proven on Histopathology	Sonography			
		True Positive	True Negative	False Positive	False Negative
Benign	18	15	19	3	2
Malignant	21	19	15	2	3

Discussion

Many breastfeeding women experience breast symptoms including pain, tenderness, firmness, and palpable lumps. Thanks to the increasing breast cancer awareness, these patients are usually referred for further examination. The lactating breast is under the influence of circulating hormones which lead to glandular proliferation, ductal dilatation, and stromal involution. Hence, the physical examination of the lactating breasts is difficult, and radiologic evaluation is usually necessary. [9] It has been suggested that, regardless of the lactational or gestational status, for symptomatic women younger than 30 years of age, US should be the first-line imaging and mammography should be saved for patients with indeterminate or suspicious lesions on US scans. US is quite successful in demonstrating true masses as well as normal breast parenchyma which may show palpable nodularity during lactation. When necessary, mammography can be performed just after breastfeeding to avoid high-density parenchyma related to retained milk products. [10] Magnetic resonance imaging findings may be altered due to physiologic changes but remains an efficient technique for the detection and characterization of breast lesions during lactation. [11]

In our study, the enrolled women were with a wide age range between 18 and 70 years. The mean age was 48.3 years old. This study is similar to study by Cacala SR et al, where mean age of the women was 47 years, with a range of 18-88 years and near similar to studies of Brennan M et al and Ayyappan AP MS et al, where age range of 14-70 and mean age of 41 years were reported. [12-14] In our study 19 patients were breast lesions occurred in the age group of 40-60 years, 6 were younger than 40 years and 14 were between 61-70 years. Most of the patients with benign (37.31%) according to BI-RADS assessment were within the age range of 40-49 years. This finding is in agreement with the results of Baker TP et al, where they found patients with malignancy to be from the 4th decade of life. [15] In our study 13 women were fibroadenoma was the commonest. Fibrocystic changes and inflammatory changes were the main categories of other lumps. According to Two studies had a higher rate compared to our study. In the young age group 73% had fibroadenoma. [16] These results may be explained by differences in ethnicity in these studies. In our study 9 patients of benign lesions were fibrocystic changes which was comparable to the result of Litton JK et al who reported 14%. [17] Four patients had mastitis and most of them were above 30 years old. This incidence

was lower compared to the other studies except for Kang YD et al, who noted mastitis in only 2.5% of their patients. [18-20] Other benign lumps were rare in our and other studies. Rare entities included phyllodes tumor, galactocele, lipoma and sclerosing adenosis. Therefore, it was difficult to compare these results. In general, in western countries, 90% of lumps are benign. The largest number of women with benign condition present with fibrocystic changes (38%), cysts (15%), fibroadenoma (13%), inflammatory (8%) and few other entities. [21]The mean age of patients in the two benign and malignant groups in our study were 42.37 years and 53 years respectively, which is similar to study by Tirada N et al who have reported significantly older age among the malignant breast lesions compared with the benign group. [22]

Limitation of the study

This study conducted at tertiary care hospital, only patients attending the hospital were enrolled. It is not possible to know how many patients with a lump did not attend the hospital. Although the probability for a woman in India to develop breast cancer is not known, it would be helpful to have larger studies to determine the extent of the disease especially in young age groups and to establish other risk factors.

Conclusion

Substantial physiological changes during pregnancy and lactation make it challenging to evaluate patients presenting with a breast problem. Most findings in pregnant and lactating patients are benign. Ultrasound is the first-line imaging modality for all pregnant women and for lactating women less than 30 years of age pregnant, or lactating and for guiding interventional breast procedures. US has a significant role in the postoperative assessment of patients with breast cancer. It is helpful in evaluating postoperative recurrent breast masses and postsurgical complications, such as infection and fat necrosis, as well as exclusion of recurrent disease.

Reference

1. Teberian I, Bhimani C, Sciotto M, Wilkes A, Germaine P. Breast masses in pregnancy and lactation. *J Am Osteopath Coll Radiol*. 2019;8:5-16.
2. Robbins J, Jeffries D, Roubidoux M, Helvie M. Accuracy of diagnostic mammography and breast ultrasound during pregnancy and lactation. *AJR Am J Roentgenol*. 2011;196(3):716-722.
3. Joshi S, Dialani V, Marotti J, Mehta TS, Slanetz PJ. Breast disease in the pregnant and lactating patient: radiological-pathological correlation. *Insights Imaging* 2013;4:527-538.
4. Taylor D, Lazberger J, Ives A, Wylie E, Saunders C. Reducing delay in the diagnosis of pregnancy-associated breast cancer: how imaging can help us. *J Med Imaging Radiat Oncol*. 2011;55(1):33-42.
5. Vashi R, Hooley R, Butler R, Geisel J, Philpotts L. Breast imaging of the pregnant and lactating patient: imaging modalities and pregnancy-associated breast cancer. *AJR Am J Roentgenol* 2013;200:321-328.
6. Son EJ, Oh KK, Kim EK. Pregnancy-associated breast disease: radiologic features and diagnostic dilemmas. *Yonsei Med J*. 2006;47(1):34-42.
7. Ahn BY, Kim HH, Moon WK, Pisano ED, Kim HS, Cha ES, et al. Pregnancy- and lactation-associated breast cancer: mammographic and sonographic findings. *J Ultrasound Med* 2003; 22:491-497.
8. Expert Panel on Breast Imaging, diFlorio-Alexander RM, Slanetz PJ, Moy L, Baron P, Didwania AD, et al. ACR Appropriateness Criteria. Breast Imaging of Pregnant and Lactating Women. *J Am Coll Radiol* 2018;15:S263-S275.
9. Whang IY, Lee J, Kim KT. Galactocele as a changing axillary lump in a pregnant woman. *Arch Gynecol Obstet*. 2007;276(4):379-382.
10. Canoy JM, Mitchell GS, Unold D, Miller V. A radiologic review of common breast disorders in pregnancy and the perinatal period. *Semin Ultrasound CT MR* 2012;33:78-85.
11. Oh YJ, Choi SH, Chung SY, Yang I, Woo JY, Lee MJ. Spontaneously infarcted fibroadenoma mimicking breast cancer. *J Ultrasound Med*. 2009; 28(10):1421-1423.
12. Cacala SR. Breast conditions during pregnancy and lactation: an understanding of unique breast conditions associated with pregnancy and lactation is essential for evaluation and management of breast problems in pregnant or lactating women. *Contin Med Educ* 2010;28:508-512.
13. Brennan M, Houssami N, French J. Management of benign breast conditions. Part 2—breast lumps and lesions. *AustFam Physician*. 2005;34(4):253-255.
14. Ayyappan AP, Kulkarni S, Crystal P. Pregnancy-associated breast cancer: spectrum of imaging appearances. *Br J Radiol*. 2010;83(990):529-534.

15. Baker TP, Lenert JT, Parker J, Kemp B, Kushwaha A, Evans G, Hunt KK. Lactating adenoma: a diagnosis of exclusion. *Breast J.* 2001;7(5):354-357.
16. Saunders C, Taylor D, Ives A. The role of breast imaging during pregnancy and lactation in the diagnosis of breast malignancy. *Radiographer* 2012;59:119-123.
17. Litton JK, Theriault RL, Gonzalez-Angulo AM. Breast cancer diagnosis during pregnancy. *Womens Health (LondEngl).* 2009;5(3):243-249.
18. Kang YD, Kim YM. Comparison of needle aspiration and vacuumassisted biopsy in the ultrasound-guided drainage of lactational breast abscesses. *Ultrasonography* 2016;35:148-152.
19. Barnes DM, Newman LA. Pregnancy-associated breast cancer: a literature review. *Surg Clin North Am.* 2007;87(2):417-430.
20. Tremblay E, Therasse E, Thomassin-Naggara I, Trop I. Quality initiatives: guidelines for use of medical imaging during pregnancy and lactation. *Radiographics* 2012;32:897-911.
21. Liberman L, Giess CS, Dershaw DD, Deutch BM, Petrek JA. Imaging of pregnancy-associated breast cancer. *Radiology* 1994;191:245-248.
22. Tirada N, Dreizin D, Khati NJ, Akin EA, Zeman RK. Imaging pregnant and lactating patients. *Radiographics* 2015;35:1751-1765.

Conflict of Interest: Nil

Source of support:Nil