

## Retrospective study to identify the fate of the third molar along the line of fracture in mandibular angle fracture

Rashmi\*

Assistant Professor, Department of Maxillofacial Surgery, Narayan Medical College and Hospital, Sasaram, Bihar, India

Received: 17-10-2020 / Revised: 29-12-2020 / Accepted: 07-01-2021

### Abstract

**Aim:** The aim of this retrospective study was to identify the fate of the third molar along the line of fracture in mandibular angle fracture. **Materials and Methods:** A Retrospective study was conducted in the Department of Maxillofacial surgery, Narayan Medical College and Hospital, Sasaram, Bihar, India for 1 year Total 80 cases were enrolled for the study, divided into two groups – group 1 in which third molar was retained and group 2 in which third molar was extracted. The cause for removal included development of pain, redness, discharge indicating periodontal problems including mobility and periapical lesions. Keeping the progress of healing of the bone in mind these teeth were extracted. **Result:** The mean age group of the population of the study was 32.54 (18 to 55 years), out of which majority of the cases 87.5% (70) were male patients and 12.5% (10) were female patients. There were a total of 80 patients with mandibular angle fracture, 35% (n=28) the third molar was removed, while in 65% (n=52) retained. The etiology of the cases was attributed to RTA and assault, the former being in majority of cases, i.e. 90% and 10% respectively. The side of the angle fracture in this study were almost similar, incidence of left side fracture being 56.25% and the right side fracture being 43.75%. In 43.75% (35) cases the third molar was completely erupted and 56.25% (45) were impacted third molars. In the impacted molars 51.11% were mesioangular, 37.78% were vertical and 11.11% were distoangular. At the end of the 4 month, 7 cases showed signs of infection in the retained group due to which extraction of the third molar was carried out under local anesthesia following aseptic precautions. There were no re fractures during extraction. Out of the 80 cases included in this study 11 cases underwent implant removal. 7 case in retained group and 4 in the removal group. P value of 0.11 was noted and a Chi square value of 1.77. **Conclusion:** We conclude that the partially impacted teeth are best to be removed during the procedure for better outcomes provided the fractured segments stability is maintained.

**Keywords:** Mandibular angle, Third molar, Tooth in line of fracture.

This is an Open Access article that uses a fund-ing model which does not charge readers or their institutions for access and distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>) and the Budapest Open Access Initiative (<http://www.budapestopenaccessinitiative.org/read>), which permit unrestricted use, distribution, and reproduction in any medium, provided original work is properly credited.

### Introduction

The mandible is the area of the face with major incidence of fracture. Its prominence and position in the skeletal face predispose to frequent traumas. Some studies have demonstrated that it is really common to observe teeth in the line of fractures.[1] Others authors mentioned that the presence of the teeth can be one of the determinant factor of the fracture location. The management of teeth in the line of fracture had changed within the past years. In the past, it was thought that teeth in the line of fractures should be immediately removed.[2,3] Although recent studies, support the vision that noninfected teeth in the line of fracture can be preserved.<sup>1</sup> The maintenance of these teeth can favor the treatment in some cases; therefore they contribute for the stability of the fracture. Its removal can be harmful, once that can diminish the contact between fragments, cause additional

trauma to the region, increase the risk of contamination of the fracture through the empty alveolus, convert a closed fracture into an open fracture and cause the loss of the bony bunch in the zone of tension. A retained tooth is when, after normal eruption, is still covered by bone and/or soft tissue.[4] This can occur due to: early loss of deciduous teeth, dental anomalies, poor positioning of the dental germ or of adjacent teeth, lack of space for eruption, permanence of deciduous teeth, trauma, impacted hard, soft, or both tissues, supernumerary teeth, odontogenic cysts, and/or tumors.[5,6] Verri et al[7] indicated that lower third molars are the teeth that most remain impacted, followed by upper third molars, upper canines, and supernumerary teeth. These data were confirmed by other authors.[4,5] Retained third molars can be classified according to the angle of the third molars' long axis compared with the adjacent second molars.[8] The vertical position is the most common, followed by mesial or mesioangulated,[9] but and Farish and Bouloux[10] ensured that the mesioangulated lower third molars are the most common position, followed by the vertical. Thus this study aimed to determine, whether the tooth in line of fracture predisposed to infection which in turn may lead to retrieval of implants.

\*Correspondence

Dr. Rashmi

Assistant Professor, Department of Maxillofacial Surgery, Narayan Medical College and Hospital, Sasaram, Bihar, India  
E-mail: [gdcgautam@gamil.com](mailto:gdcgautam@gamil.com)

## Material and Methods

The present Retrospective study was conducted in the Department of Maxillofacial surgery Narayan Medical College and Hospital, Sasaram, Bihar, India for 1 year.

### Inclusion criteria

- Patients with angle fracture that required open reduction and internal fixation,
- Age between 18-55years.

### Exclusion criteria

- Patients with pre-existing medical conditions
- Infected fracture site
- Patients who were treated by closed reduction
- Patients having less than 6 months follow up

## Methodology

Intraoral vestibular incision was used to approach the fracture, anatomic reduction was obtained and plating was done. The third molar was removed when the teeth were fractured, pre-existing pericoronal/periodontal infection, dental caries, tooth mobility, exposure or involvement of the apical half or more of the root, and third molar does not compromise the reduction of bone fragments. For the purpose of this study, postoperative infection was defined as that has a purulent discharge requiring surgical intervention including removal of plates. For all the cases semi-rigid fixation was done with plates and screws after fracture reduction, standard analgesics and antibiotic coverage were given. Total 80 cases were enrolled for the study, divided into two groups – group 1 in which third molar was retained and group 2 in which third molar was extracted. The cause for removal included development of pain, redness, discharge indicating periodontal problems including mobility and periapical lesions. Keeping the progress of healing of the bone in mind these teeth were extracted. At the 3rd month follow up 7 teeth which were initially retained were extracted when signs of infection were first seen. At the 6th month follow up the implants were removed along with extraction of the teeth, in

the retained group i.e 7 in number and 4 cases of implant removal in the removed group.

## Statistical analysis

The data were evaluated using SPSS 19.0. Significant differences between the various approaches were identified using chi-square test. The level of confidence interval and p-value were set at 95% and 5%

## Results

The mean age group of the population of the study was 32.54 (18 to 55 years), out of which majority of the cases 87.5% (70) were male patients and 12.5% (10) were female patients (Table 1). There were a total of 80 patients with mandibular angle fracture who underwent open reduction and internal fixation. 35% (n=28) the third molar was removed, while in 65% (n=52) retained (Table 2). The etiology of the cases was attributed to RTA and assault, the former being in majority of cases, i.e. 90% and 10% respectively. The side of the angle fracture in this study were almost similar, incidence of left side fracture being 56.25% and the right side fracture being 43.75%. In 43.75% (35) cases the third molar was completely erupted and 56.25% (45) were impacted third molars. In the impacted molars 51.11% were mesioangular, 37.78% were vertical and 11.11% were distoangular (Table 3). At 4 months follow up, based on the signs of infection it was decided to extract the third molars. The signs and symptoms considered were –pain, redness or discharge at the third molar site. At the end of the 4 month, 7 cases showed signs of infection in the retained group due to which extraction of the third molar was carried out under local anesthesia following aseptic precautions. There were no re fractures during extraction. Out of the 80 cases included in this study 11 cases underwent implant removal. 7 in case in retained group and 4 in the removal group. In the retained group, extraction of the third molar was carried out along with removal of the implant. A p value of 0.11 was noted and a Chi square value of 1.77 (Table 4).

**Table 1: Demographic Profile**

Gender	N=80	%
Male	70	87.5
Female	10	12.5
<b>Age</b>		
Below 20	6	7.5
20-40	62	77.5
40-60	12	15
<b>RTA</b>	70	90

**Table 2: Distribution of the subjects based on retainment or removal of third molar**

Retainment or removal	N=80	%
3 <sup>rd</sup> molar Removed Group	28	35
3 <sup>rd</sup> molar Retained Group	52	65
Total	80	100.0

**Table 3: Cross-tabulation of 3<sup>rd</sup> molar impaction and type of impaction**

Type of impaction		3 <sup>rd</sup> molar impaction		Total
		Complete	Partial	
Not applicable	Count	35	0	35
	Percent	100.0%	0.0%	43.75%
Disto-angular	Count	0	5	5
	Percent	0.0%	11.11%	6.25%
Mesio-angular	Count	0	23	23
	Percent	0.0%	51.11%	28.75%
Vertical	Count	0	17	17
	Percent	0.0%	37.78%	21.25%
Total	Count	35	45	80
	Percent	100.0%	100.0%	100.0%

**Table 4: Cross-tabulation of 3<sup>rd</sup> molar retainment and implant retrieval**

Implant retrieval		3 <sup>rd</sup> molar retainment		Total
		Retained	Removed	
Retained	Count	21	48	69
	Percent	75%	92.31%	86.25%
Retrieval	Count	7	4	11
	Percent	25%	7.69%	13.75%
Total	Count	28	52	80
	Percent	100.0%	100.0%	100.0%

Chi-square value- 1.77, P value- 0.11

## Discussion

This has always been a question of debate and the risk pertaining to retaining or removing the tooth has been varyingly assessed in literature ever since evolution of open reduction and fixation for maxillo facial fractures were introduced. In the present study, angle fracture was observed in the age group ranging from 18 to 55 years and the mean age was 32.54 years. Based on age the patients were classified into three categories i.e. younger age group below 20 years years, middle aged group 20 to 40 years, and older age group - above 40 years. Out of the 80 patients included in the study, 6 belonged to the young group, 62 to the middle age group and 12 to the old age group, indicating that majority of the angle fractures occurred in middle age group, and road traffic accidents being the most common cause of it. This result was in consistent with the results of the study conducted by Sakr et al, who reported that incidents of angle fracture between 20-29 years are higher. The reason is due to the fact that a high incidence of un-erupted third molars is seen in this age group.[11] Our study consisted of 87.5% of male patients and 12.5% of female patients. This observation was in agreement with studies conducted by Dongas et.al and Mahesh Kumar et al who reported male predominance in angle fractures due to the fact that they are more exposed to

the risk factors for facial trauma as they are prone to get involved in violent conduct, indulging in reckless driving, exhibiting physical aggression and engaging in contact sports.[12,13] The majority of the cases had an etiology of road traffic accidents i.e. 90 and 10% of cases had an etiology of assault. This result was consistent with the study conducted by Ugboko et who had observed that road traffic accidents were the main cause of mandibular angle fractures. This is attributed to multiple reasons, but the main reason being lack of road safety awareness, violation of traffic rules like over-speeding and not using helmet, use of alcohol or other intoxicating agents.[14] We found 45 (56.25%) cases of mandibular angle fracture on the left side as compared to 35 (43.37%) on the right side. This was in agreement with the study findings of Inaoka et al., where they proved left side had more angle fractures than the right side. However, the side did not present a significant relationship with angle fracture. The site of impact is usually restricted to the side of fall. If the impact is of a high velocity, then a direct fracture at the point of application will occur. If the impact is of a low velocity, the bow will transfer to the contralateral side, causing an indirect fracture.[9] In case of assaults, considering the predominance of the right-handed people, the victim will be facing the opposite direction and hence the site of fracture is to the side of impact. In our study we noted that

all the assault cases had an angle fracture on the left side. In our study 43.75% of the cases had their third molar completely erupted whereas as 56.25% of the cases exhibited impaction of the third molar due to the fact that majority of the cases belonged to the young age group. Among the impacted cases, it was noted that mesioangular impaction was the most common type of impaction this was in agreement to the study findings of Fuselier et al.[15] it was attributed that mesioangular impacted teeth are more prone to angle fracture as the root is directed towards the angle of mandible, which may act as a wedge splitting the mandibular angle, because of which the injury forces are redirected towards the mandibular angle, and decreased amount of bone in that area increases the risk of angle fracture. Mandibular angle fractures observed along with other impaction positions of third molars in decreasing order were: Vertical, horizontal, and distoangular. The type of impacted teeth did not have a role in deciding whether the tooth needed to be removed or retained intra-operatively. In the post-operative follow – up period it was noted that signs of infection which led to the removal of the impacted teeth were noted more in partially impacted cases. The study conducted by Balaji et al was in agreement to our stud, this was simply because of the position of the tooth which makes it an area for harboring debris and pathogen which in-turn led to periodontal infection.[16] In a recent systematic review by Bobrowski et al, of the 1542 cases, tooth was removed in 788 (51.1%). During the follow-up period infection occurred in 84 cases (10.66%). On other hand, 84 cases out of 754 in the retained group showed signs of infection. This had no statistical significance. Thus the study was concluded by saying that retaining or removing the third molar did not have a significant effect on infection.[17] In an article by Ellis et al, Muller had recommended that multi-rooted tooth in the line of fracture be always removed.[18] In another similar study with the same sample size conducted by Lim et al, 49 patients had third molars in the line of fracture. The third molars were retained in 39 cases and the third molars were extracted in the rest of the cases. It was noted that several patients in the retained group exhibited post op infections, nerve paresthesia, temporomandibular disorders and also change of occlusion. Whereas in case of the group in which the third molars were extracted, they noticed that the patients presented with only nerve injury. However this study also did not yield a statistically significant value.[19] In our study we did not encounter any TMD or nerve injury cases, although post-operative infection was noted. In a study conducted by Kahnberg and Ridell it was found that the teeth which were retained along the fracture line resulted in satisfactory healing, which was around 59%.[20] This was later supported by works of Macan et al.[21] Other teeth have relatively better access and survival rate with adjuvant treatments like root canal therapy while the third molar would lack the same. Also, this study proves that fully impacted third molar teeth when removed did not cause any further infection, while the partially impacted teeth which were left behind, proceeded to infection and subsequent loss of teeth. In our study the difference in survival of third molar was not statistically significant between right and left side. The partially impacted

teeth, due to its position would harbor more debris and pathogens contributing to poor periodontal health. Although this finding did not yield a statistically significant it was what we inferred from our study.

### Conclusion

Retaining the third molar has an increased chance of post-operative infections it is not statistically significant. Other reasons also lead to the post-operative infections. We conclude that the partially impacted teeth are best to be removed during the procedure for better outcomes provided the fractured segments stability is maintained.

### References

1. Neal DC, Wagner WF, Alpert B. Morbidity associated with teeth in the line of mandibular fractures. *J Oral Surg* 1978; 36:859-862.
2. Amaratunga NA. A comparative study of the clinical aspects of edentulous and dentulous mandibular fractures. *J Oral Maxillofac Surg* 1988; 46:3-5.
3. Bradley RL. Treatment of fractured mandible. *Amer Surg* 1965; 31: 289-290.
4. Peterson LJ, Ellis E III, Hupp JR, Tucker M. *Contemporary Oral & Maxillofacial Surgery*. Saint Louis: Mosby; 1993:225–261; 587– 610.
5. Damante JH, Freitas JAS, Tavano O, Alvares LC. Interpretação radiográfica. In: Alvares LC, Tavano O, eds. *Curso de Radiologia em Odontologia*. São Paulo: Editora Santos; 2009:129–218.
6. Al-Khateeb TH, Bataineh AB. Pathology associated with impacted mandibular third molars in a group of Jordanians. *J Oral Maxillofac Surg* 2006;64(11):1598–1602.
7. Verri VA, Oliveira MA, Grandini AS, et al. Estudo clínico-radiográfico da incidência dos dentes inclusos em 3.000 indivíduos. *Rev Assoc Paul Cirurg Dent* 1973;27(5):274–279.
8. Marciani RD. Third molar removal: an overview of indications, imaging, evaluation, and assessment of risk. *Oral Maxillofac Surg Clin North Am* 2007;19(1):1–13.
9. Inaoka SD, Carneiro SCAS, Vasconcelos BCE, Leal J, Porto GG. Relationship between mandibular fracture and impacted lower third molar. *Med Oral Patol Oral Cir Bucal* 2009;14(7): E349–E354.
10. Farish SE, Bouloux GF. General technique of third molar removal. *Oral Maxillofac Surg Clin North Am* 2007;19(1):23–43.
11. Sakr K, Farag IA, Zeitoun IM. Review of 509 mandibular fractures treated at the University Hospital, Alexandria, Egypt. *Br J Oral Maxillofac Surg*. 2006;44(2):107-11.
12. Dongas P, Hall GM. Mandibular fracture patterns in Tasmania, Australia. *Aust Dent J*. 2002;(2):131-7.
13. Subbaiah MK, Ponnuswamy I, David M. Relationship between mandibular angle fracture and state of eruption of mandibular third molar: A digital radiographic study. *J Indian Acad Oral Med Radiol*. 2015;27(1):35.

14. Ugboko VI, Oginni FO, Owotade FJ. An investigation into the relationships between mandibular third molars and angle fractures in Nigerians. *Br J Oral Maxillofac Surg.* 2000;38(5):427-9.
15. Fuselier JC, Ellis EE, Dodson TB. Do mandibular third molars alter the risk of angle fracture? *J Oral Maxillofac Surg.* 2002;60(5):514-8.
16. Balaji P, Balaji SM. Fate of third molar in line of mandibular angle fracture - Retrospective study. *Indian J Dent Res.* 2015;26(3):262-6.
17. Bobrowski AN, Sonogo CL, Chagas OL. Postoperative infection associated with mandibular angle fracture treatment in the presence of teeth on the fracture line: A systematic review and meta-analysis. *Int J Oral Maxillofac Surg.* 2013;42(9):1041-8.
18. Ellis E. Outcomes of patients with teeth in the line of mandibular angle fractures treated with stable internal fixation. *J Oral Maxillofac Surg.* 2002;60(8 SUPPL.1):863-5.
19. Lim HY, Jung TY, Park SJ. Evaluation of postoperative complications according to treatment of third molars in mandibular angle fracture. *J Korean Assoc Oral Maxillofac Surg.* 2017;43(1):37-41.
20. Kahnberg KE, Ridell A. Prognosis of teeth involved in the line-of mandibular fractures. *Int J Oral Surg.* 1979;8(3):163- 72.
21. Kamboozia AH, Punnia-Moorthy A. The fate of teeth in mandibular fracture lines. *Int J Oral Maxillofac Surg.* 1993;22(2):97-101.

**Conflict of Interest: Nil**

**Source of support:Nil**