

To evaluate the various indications and the outcomes of tracheostomy in ICU; an observational study

Indrajeet Kumar¹, Birendra Kumar^{2*}

¹Assistant Professor, Department of ENT, Nalanda Medical College and Hospital, Patna, Bihar, India

²Associate Professor & HOD, Department of ENT, Nalanda Medical College and Hospital, Patna, Bihar, India

Received: 12-10-2020 / Revised: 29-12-2020 / Accepted: 8-01-2021

Abstract

Objective: to study the various underlying disease aspects of indications and the outcomes of tracheostomy in ICU at Nalanda Medical College and Hospital, Patna, Bihar, India. **Material and methods:** This cross-sectional study was done in ICU, Department of ENT Nalanda Medical College and Hospital, Patna, Bihar, India for 9 months. Total 110 patients who underwent elective open tracheostomy in ICU by ENT surgeons for various indications were included in this study. Patients details (age and sex), complete clinical history pertaining to the cause of prolonged intubation/ indication of tracheostomy, timing and complications was compiled and analyzed. **Results:** Out of 110 patients 75 (68.19%) were male and 35 (31.81%) were female and male to female ratio was 2.14:1. The age ranges of patients from 12-72 years and highest frequency were 20-30 years age group 35 (31.81%) followed by 25(22.75%) were from 40-50 years age group. Out of 110 cases head injury with h/o road traffic accident (RTA) was the frequent indication 30 (27.27%) and postoperative case of intracranial space occupying lesion (ICSOL) was the second most indication 28 (25.45%). Out of 110 cases over all complications was 11 (10%) and most common complication was surgical emphysema 4 (3.36%) followed by hemorrhage 3 (2.72%), tube displacement 2(1.81%) and wound infection 2 (1.81%). **Conclusion:** Tracheostomy in ICU is an important and safe procedure if prolonged endotracheal intubation is advised for varying underlying causes.

Keywords: Tracheostomy, Intensive care unit.

This is an Open Access article that uses a fund-ing model which does not charge readers or their institutions for access and distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>) and the Budapest Open Access Initiative (<http://www.budapestopenaccessinitiative.org/read>), which permit unrestricted use, distribution, and reproduction in any medium, provided original work is properly credited.

Introduction

Tracheostomy is described as the creation of a stoma at the skin surface which leads into the trachea which was at the earliest performed in ancient Egypt and is also included in many ancient medical texts like The Rig Veda (2000 to 1000 BCE)[1,2]. Tracheostomy is performed in about 24% of all patients in intensive care units (ICU)[3,4]. Tracheostomy has many advantages over endotracheal intubation in ICU setting including protection of the larynx and the upper airway from prolonged intubation which may lead to tracheal stenosis [5], improved patient comfort, less requirement for sedation faster weaning leading to reduce ICU and hospital stay and reduced incidence of ventilator associated pneumonia if done early[6,7].

Indications are mainly prolonged intubation for various conditions, acute or chronic neuromuscular diseases, poor cardio- respiratory reserve, brain injury and upper airway obstruction[8]. While the timing of tracheostomy differs for these indications and its recommended for consideration only if extubation did not occur by 21 days in prolonged cases⁸ but in selected patients with severe multi-trauma and/or head injury with low Glasgow coma score, tracheostomy at the earliest, within 3–4 days of intubation is advocated[9]. Tracheostomy has been considered a safe procedure in ICU but has been found to lead to life-threatening complications intra and post operatively like hypoxia, cardiac arrest, injury to structures immediately adjacent to the trachea, pneumothorax, haemothorax, incision site bleeding and stoma infection[10]. Techniques of tracheostomy include open surgical technique, earlier performed routinely and percutaneous dilatational tracheostomy (PDT) advocated by Ciaglia[11], which is commonly performed in recent days in ICU because of its various

*Correspondence

Dr. Birendra Kumar

Associate Professor & HOD, Department of ENT, Nalanda Medical College and Hospital, Patna, Bihar, India

E-mail: dr.birendra1954@gmail.com

advantages over the former, but PDT is not routinely practiced in our institution. Although recent studies have suggested that tracheostomy can be a safe procedure in the ICU, tracheostomy has also been found to lead to life-threatening complications like hypoxia, cardiac arrest, injury to structures immediately adjacent to the trachea, pneumothorax and haemothorax. Many critically ill patient's families have been hesitant in authorizing tracheostomy because of cosmetic issues and speech problems[12,13]. In the recent years more and more airway problems are managed with endotracheal intubation or percutaneous endoscopically guided tracheostomy. But in many countries percutaneous endoscopically guided tracheostomy is not yet routinely practiced, conventional tracheostomy is practiced in vast majority of cases in ICU. The objective of present study is to study the various underlying disease aspects of indications and the outcomes of tracheostomy in ICU at Nalanda Medical College and Hospital, Patna, Bihar, India.

Material and methods

This cross-sectional study was done in ICU, Department of ENT Nalanda Medical College and Hospital, Patna, Bihar, India for 9 months among the patients having tracheostomy .

Inclusion criteria

- All patients who underwent tracheostomy
- Patient already intubated in ICU assumed to require prolonged intubation period.

- Patient with pre-operative planned tracheostomy and post operatively stay in ICU.

Exclusion criteria

- Patients not willing for the study
- Not available for follow up.
- Patients who have undergone tracheostomy at other hospital

Methodology

Total 110 patients who underwent elective open tracheostomy in ICU by ENT surgeons for various indications were include in this study. Patients details (age and sex), complete clinical history pertaining to the cause of prolong intubation/ indication of tracheostomy, timing and complications was compiled and analyzed.

Results

Out of 110 patients 75 (68.19%) were male and 35 (31.81%) were female and male to female ratio was 2.14:1 (Table 2). the age ranges of patients from 12-72 years and highest frequency were 20-30 years age group 35 (31.81%) followed by 25(22.75%) were from 40-50 years age group.table 1 Out of 110 cases head injury with h/o road traffic accident (RTA) was the frequent indication 30 (27.27%) and postoperative case of intracranial space occupying lesion (ICSOL) was the second most indication 28 (25.45%) (Table 2).

Table 1: Demographic Profile of Patients

Gender	Number of patients=110	%
Male	75	68.19
Female	35	31.81
Age in years		
Below 20 year	8	7.27
20-30	35	31.81
30-40	20	18.18
40-50	25	22.72
50-60	16	14.54
Above 60	24	21.82

Table 2: Indications of tracheostomy (n=110)

Indications	Number of patients=110	%
Head injury with h/o RTA	30	27.27
Post-operative case of ICSOL	28	25.45
Guillain-Barre syndrome	23	20.90
CVA	12	10.60
Maxillo-facial trauma	5	4.54
RTA with spinal cord injury	7	6.36
Post-operative pneumonia	5	4.54

Out of 110 cases over all complications was 11 (10%) and most common complication was surgical emphysema 4 (3.36%) followed by hemorrhage 3 (2.72%), tube displacement 2(1.81%) and wound infection 2 (1.81%) (Table 3).

Table 3: Complications of tracheostomy (n=110)

Complications of tracheostomy	Number of patients=11	%
Surgical emphysema	4	3.36
Haemorrhage	3	2.72
Tube displacement	2	1.81
Wound infection	2	1.81

Discussion

Tracheostomy is a common procedure done in ICU having its own merit and demerits but not always without complications. It is one of the life-saving operations. In this study 110 cases of tracheostomy in the ICU were studied and the patient demographics, indications, complications and outcomes were analyzed and compare with similar study. In this study age ranges of patients from 12-72 years and highest frequency were 20-30 years age group 35 (31.81%) followed by 25(22.75%) were from 40-50 years age group. One study done by Chandra et al showed maximum age group between 21-30 years age group 28% followed by 21% were from 41-50 years age group which is similar to our study[14]. In our study sex distribution among the 110 cases of tracheostomy in ICU showed 75 (68.19%) were male and 35 (31.81%) were female and male to female ratio was 2.14:1. Study done by Mahmud et al male patient were 65% and female patients were 35%, male to female ratio 1.8:1 which is similar to our study[15]. Another study done by Chandra et al male patients were 61% and female patients were 39% which nearer to our study[14]. Study done by Perfeito et al showed male to female ratio was found 1.8:1 which is similar to our study[16]. In this study, the commonest indications of tracheostomy in ICU were head injury with h/o RTA was the frequent indication 30 (27.27%) followed by post-operative case of ICSOL was 28 (25.45%). One study done by Chowdhury et al showed head injury with H/O RTA 26.67% followed by post-operative case of ICSOL 26.67% which is similar to our study[17]. Another study done by Mahmud et al showed head injury with RTA 27.5% followed by post-operative case of ICSOL were 25% which is nearer to our study[15]. The incidence of RTA is very high in our country due to overloaded or unroadworthy vehicles, lack of awareness of safe road use, poor traffic management and law enforcement and poor driver training[18]. In our study, the rate of complications of tracheostomy in ICU was 11 (10%). Study done by Mahmud et al was 10%, Perfeito et al was 8.7% and Chowdhury et al was 10% which is nearer to our study[15-17]. In our study, the most common complication of tracheostomy in ICU was surgical

emphysema 4 (3.36%) followed by hemorrhage 3 (2.72%), tube displacement 2(1.81%), wound infection 2 (1.81%). Study done by Mahmud et al and Chowdhury et al showed similar result[15-17]. Another study done by Perfeito et al showed early complication was bleeding while late complication was wound infection in 2.73% which is near to our study[16]. Another study done by Datta et al showed 2% cases had wound infection which is nearer to our study[19]. Study done by Rahman et al and Ahmed et al showed similar study about our complications[20,21]. In our study most common complication was surgical emphysema 4 (3.36%) managed by removal of tight suture, hemorrhage 3 (2.72%) control by intraoperative pressure over bleeders and ligation, wound infection 2 (1.81%) was treated by regular dressing of wound and appropriate antibiotics. Study done Mahmud et al and Chandra et al showed similar result[14,15]. Complications of tracheostomy has been extensively studied and found to be decreased with improvements in operative skill and advancements in ICU[22]. Patents and caregiver education prior to performing elective tracheostomy and during discharge will help to improve patient outcomes and decrease complications related to tracheostomy tube. Complication rates associate with tracheostomy can be prevented by use of non-metallic tube, good surgical technique and meticulous post-operative care[23]. In our study, incidence of dysphagia, aspiration, trachea-esophageal fistula, tracheo-cutaneous fistula and cardiac arrest, we found no such complications which is accordance to study done by Chowdhury et al, study done by Mahmud et al.[15,17]. No death of patient was reported during tracheostomy in our study which is accordance to study done by Chandra et al.[14] The reason of complication in our study may be due to possibility of performing most of tracheostomy by the junior doctors which is accordance to study done by Mahmud et al.[15] In this study, regarding benefits of tracheostomy over endotracheal intubation in ICU we found that 100% patents had greater comfort. Nursing care was easier especially with respects to suctioning in 100% of patient, reduction of the length of ICU study found in all cases. Better oral and airway care was possible in all

cases. This study was accordance to study done by Mahmud et al and Perfeilo et al.[15,16]

Conclusion

Tracheostomy in ICU is an important and safe procedure if prolonged endotracheal intubation is advised for varying underlying causes.

References

1. Pracy P. Tracheostomy. In: Gleeson M et al eds Scott Browns Otolaryngology and Head-Neck Surgery, London: Hodder Arnold; 2008; Vol1: p-353.
2. Frost EAM. Tracing the tracheostomy. *Annals of Otolaryngology*. 1976; 85: 618–24.
3. Upadhyay A, Maurer J, Turner J, Tiszenvel H, Rosengart T, Elective bedside Tracheostomy in the intensive care unit. *J Am Coll Surg*. 1996; 183(1): 51-5.
4. Esteban A, Anzueto A, Alia I, Gordo F, Apezteguia C, Palizas F, et al.: How is mechanical ventilation employed in the intensive care unit? An international utilization review. *Am J Respir Crit Care Med* 2000, 161; 1450-1458.
5. Blot F, Similowski T, Trouillet JL, Chardon P, Korach JM, Costa MA, et al. Early tracheotomy versus prolonged endotracheal intubation in unselected severely ill ICU patients. *Intensive Care Med* 2008; 34(10): 1779-1787.
6. Nieszkowska A, Combes A, Luyt CE, Ksibi H, Trouillet JL, Gilbert C, Chastre J. Impact of tracheostomy on sedative administration, sedation level, and comfort of mechanically ventilated intensive care unit patients. *Crit Care Med* 2005; 33(11): 2527-2533.
7. Griffiths J, Barber VS, Morgan L, Young JD. Systematic review and meta-analysis of studies of the timing of tracheostomy in adult patients undergoing artificial ventilation. *BMJ* 2005; 330(7502): 1243.
8. Bary B, Bodenhan AR. Role of Tracheostomy in ICU, *Anaesthesia & Intensive care Medicine*. 2004; P-375.
9. McWhorter AJ. Tracheotomy: timing and techniques. *Curr Opin Otolaryngol Head Neck Surg* 2003; 11(6): 473-479.
10. Stock MC, Woodward CG, Shapiro BA,: Perioperative complications of elective tracheostomy in critically ill patients. *Crit Care Med* 1986, 14: 861-863.
11. Ciaglia P, Firsching R, Syniec C. Elective percutaneous dilatational tracheostomy. A new simple bedside procedure; preliminary report. *Chest* 1985; 87(6): 715-719.
12. Zeitouni AG, Kost KM: Tracheostomy: a retrospective review of 281 cases. *J Otolaryngol* 1994, 23:61-66.
13. Stock MC, Woodward CG, Shapiro BA, : Perioperative complications of elective tracheostomy in critically ill patients. *Crit Care Med* 1986, 14:861-863.
14. Shekar CY, Viswanatha B, Srinivasan SB, Joyaram RT, Vijayashree MS. tracheostomy in intensive care unit: Indications and outcomes at a teaching Hospital. *J Otolaryngology*. 2016;5(2): 28-31.
15. Mohmud M, Hossain MA, Sarkar MZ, Hossain HSM, Islam MO, Ahmed MU, et al. Tracheostomy in intensive care unit. Indication, Benefits and complications. *Bangladesh J Otorhinolaryngology*. 2015;21(1):28-32.
16. Perfeito JAJ, Forte V, camaghi M, Tamuran N. *Jornal Brasileiro de pneumologia: tracheostomy in ICU: it is worthwhile*. *Bangladesh Crit Care J*. 2007;3(6):1-5.
17. Chowdhury AA, Sultana T, Joarder AH, Tarafder KH. A comparative study of elective and emergency tracheostomy. *Bangladesh J Otorhinolaryngology*. 2008;14(2):57-62.
18. Centre for the Rehabilitation of the paralysed. Road safety in Bangladesh; 2010.
19. Datta RK, Viswanatha B, Puneet PJ, Meerin B, Kumari TLN. Tracheostomy: Our Experience Research in otolaryngology. 2015;4(2):29-33.
20. Rahman SH, Ahmed K, Khan AFM, Ahmed SU, Hanif MA, Haroon AA, et al. Study of tracheostomy in Dhaka medical College Hospital. *Bangladesh J Otorhinolaryngology*. 2001;7(2):34-40.
21. Ahmed K, Rahinan MA, Rahman SH. Complication of tracheostomy. *Bangladesh J Otorhinolaryngology*. 1998;4(1):3-6.
22. Manuel L, Mark O. Tracheal obstruction as a complication of tracheostomy tube malfunction; case report and review of the literature. *J Bronchology Interventional Pulmonology*. 2010; 77:253-7.
23. Ashwin MA, Deepa R, Balakrishnan E, Aswin M, Anupama M. Tracheostomy: a hospital based descriptive study. *Medplus Int J ENT*. 2017;1 (2):33- 9.

Conflict of Interest: Nil

Source of support:Nil